

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
28 July 2005 (28.07.2005)

PCT

(10) International Publication Number
WO 2005/067601 A2

(51) International Patent Classification: **Not classified**

(72) Inventors; and

(21) International Application Number:

PCT/US2005/000302

(75) Inventors/Applicants (for US only): **JOHNSON, Karl, F.** [US/US]; 5320 Ivy Stream Road, Haltboro, PA 19040 (US). **BEZILA, Dan** [US/US]; 181 Montgomery Drive, Harleysville, PA 19438 (US). **NGO, Winnie** [US/US]; 116 Byberry Road, Haltboro, PA 19040 (US). **HAKES, David** [US/US]; 14 Fern Avenue, Willow Grove, PA 19090 (US).

(22) International Filing Date: 6 January 2005 (06.01.2005)

(25) Filing Language:

English

(74) Agents: **BASTIAN, Kevin, L.** et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, 8th Floor, San Francisco, CA 94111 (US).

(26) Publication Language:

English

(30) Priority Data:

60/535,263

9 January 2004 (09.01.2004)

US

(81) Designated States (unless otherwise indicated, for every kind of national protection available): **AL, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,**


(71) Applicant (for all designated States except US): **NEOSE TECHNOLOGIES, INC.** [US/US]; 102 Witmer Road, Horsham, PA 19044 (US).

[Continued on next page]

(54) Title: **VECTORS FOR RECOMBINANT PROTEIN EXPRESSION IN E.COLI**



A



B

(57) Abstract: The present invention relates to methods of providing a protein product to a customer. In particular, the invention relates methods of using protein expression vectors to produce proteins to be provided to a client. The invention also provides vectors for the cloning and expression of proteins, including reagent proteins and therapeutic proteins.



PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TITLE

Vectors for Recombinant Protein Expression in E. Coli

5

BACKGROUND OF THE INVENTION

Humans have exploited the use of genetics and microorganisms for their own advantage throughout much of recorded history. Egyptians are credited with the first use of yeast to produce leavened bread sometime between 4000-2000 BC, Gregor Mendel produced peas having specific, defined, characteristics in the mid-19th century, and the Food and Drug Administration approved the first recombinant drug, human insulin, in 1982. This last feat is often considered to be the beginning of the modern biotechnology industry. Since then, transgenic plants, recombinant foods, recombinant vaccines, cancer therapeutics, recombinant antibodies, enzymes, glycosyltransferases, cytokines, coagulation factors, hormones, dermal replacements, anti-virals, and many other recombinant proteins have been developed for human use.

The nucleic acid expression vector has greatly aided in the production of recombinant proteins and therapeutics. A nucleic acid encoding a protein reagent or therapeutic protein can be cloned into an expression vector, which can be expressed in a population of eukaryotic and/or prokaryotic cells, thus producing a large amount of a recombinant protein or therapeutic. However, the yield and quality of the recombinant product depend greatly on the expression vector and microorganism used to express the vector. In addition, the use of recombinant cells can be slow and tax resources that can be otherwise used for discovery and improvement of recombinant proteins and therapeutics.

As the demand and usefulness of recombinant proteins increases, new methods are required in order to more efficiently prepare such proteins with a rapid turnaround time. Moreover, as recombinant proteins for the treatment of a variety of diseases are generated, methods to lower the cost of their production need to be

implemented so that these technologies are available to all those in need. The need to provide improved vectors for protein expression does not exist solely in the therapeutic protein arena. Rather, this need also extends to the production of proteins or reagents (e.g., enzymes) for use in the production of both protein and non-protein therapeutics.

Over the past several decades, recombinant proteins and therapeutics have proven to be the answer in treating many diseases that were not addressed using conventional, chemical therapeutics. However, recombinant technology has been hampered by inefficiency, especially in small scale situations, as well as high cost and slow turnaround time. Streamlining the expression of proteins at a lower cost with a quicker turnaround time for virtually any customer situation is needed to realize the potential of recombinant proteins as reagents and as therapeutics. The present invention meets this need.

BRIEF SUMMARY OF THE INVENTION

The invention includes a method of providing a therapeutic protein to a customer, comprising cloning a nucleic acid encoding a protein into a pCWin1 expression vector as set forth in SEQ ID NO:1, expressing a protein therefrom, and providing the protein to a customer.

In another aspect of the invention, a method of providing a therapeutic protein to a customer comprises cloning a nucleic acid encoding a protein into a pCWin2 expression vector as set forth in SEQ ID NO:2, expressing a protein therefrom, and providing the protein to a customer.

In yet another aspect of the invention, a method of providing a therapeutic protein to a customer comprises cloning a nucleic acid encoding a protein into a pCWin2-MBP expression vector as set forth in SEQ ID NO:3, expressing a protein therefrom, and providing the protein to a customer. In still another aspect, a method of providing a therapeutic protein to a customer comprises cloning a nucleic acid encoding a protein into a pCWin2-MBP-SBD (pMS₃₉) expression vector as set forth in SEQ ID NO:10, expressing a protein therefrom, and providing the protein to a customer. In yet another aspect of the invention, a method of providing a therapeutic protein to a customer comprises cloning a nucleic acid encoding a protein into a

pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector as set forth in SEQ ID NO:11, expressing a protein therefrom, and providing the protein to a customer.

In an embodiment of the invention, a pCWin2/MBP vector comprises a protease cleavage site coding sequence between the MBP coding sequence and the therapeutic protein coding sequence.

Therapeutic proteins useful in the present invention include erythropoietin, human growth hormone, granulocyte colony stimulating factor, interferons alpha, -beta, and -gamma, Factor IX, follicle stimulating hormone, interleukin-2, erythropoietin, anti-TNF-alpha, and lysosomal hydrolases such as beta-glucosidase, alpha-galactosidase-A, beta-hexosaminidase, beta-galactosidase, alpha-galactosidase, alpha-mannosidase, beta-mannosidase, alpha-L-fucosidase, beta-glucuronidase, alpha-glucosidase, alpha-N-acetylgalactosaminidase, and acid phosphatase.

In one embodiment of the invention, a method of providing a protein to a customer includes cloning a nucleic acid encoding a protein into a pCWin1 expression vector as set forth in SEQ ID NO:1, expressing a protein therefrom, and providing the protein to a customer.

In another embodiment of the invention, a method of providing a protein to a customer includes cloning a nucleic acid encoding a protein into a pCWin2 expression vector as set forth in SEQ ID NO:2, expressing a protein therefrom, and providing the protein to a customer.

In another embodiment of the invention, a method of providing a protein to a customer includes cloning a nucleic acid encoding a protein into a pCWin2/MBP expression vector as set forth in SEQ ID NO:3, expressing a protein therefrom, and providing the protein to a customer. In still another aspect, a method of providing a protein to a customer comprises cloning a nucleic acid encoding a protein into a pCWin2-MBP-SBD (pMS₃₉) expression vector as set forth in SEQ ID NO:10, expressing a protein therefrom, and providing the protein to a customer. In yet another aspect of the invention, a method of providing a protein to a customer comprises cloning a nucleic acid encoding a protein into a pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector as set forth in SEQ ID NO:11, expressing a protein therefrom, and providing the protein to a customer.

In one aspect of the invention, a protein may be a glycosyltransferase or a sugar nucleotide-generating enzyme.

In an aspect of the invention, an expression vector includes an affinity tag coding sequence. In this aspect of the invention, an affinity tag may be a histidine tag, a Factor IX tag, a glutathione-S-transferase tag, starch-binding domain and a FLAG-tag.

- 5 The invention includes a method of providing a therapeutic protein to a customer, where the method includes providing an expression vector to a protein production facility wherein a nucleic acid encoding a protein is cloned into the expression vector and the protein is expressed therefrom in the protein production facility; subsequently providing the protein to a customer. In an aspect of the
- 10 invention, the expression vector comprises a multiple-cloning region and an antibiotic resistance marker. The antibiotic resistance marker may be kanamycin, tetracycline, or chloramphenicol. In another aspect of the invention, the expression vector includes an affinity tag.

- 15 In an embodiment of the invention, a method of providing a protein to a customer includes providing a pCWin1 vector as set forth in SEQ ID NO:1 to a protein production facility, wherein a nucleic acid encoding a protein is cloned into the expression vector and the protein is expressed therefrom in the protein production facility, and the protein is provided to a customer.

- 20 In an embodiment of the invention, a method of providing a protein to a customer includes providing a pCWin2 vector as set forth in SEQ ID NO:2 to a protein production facility, wherein a nucleic acid encoding a protein is cloned into the expression vector and the protein is expressed therefrom in the protein production facility, and the protein is provided to a customer.

- 25 In an embodiment of the invention, a method of providing a protein to a customer includes providing a pCWin2/MBP vector as set forth in SEQ ID NO:3 to a protein production facility, wherein a nucleic acid encoding a protein is cloned into the expression vector and the protein is expressed therefrom in the protein production facility, and the protein is provided to a customer. In another embodiment of the invention, a method of providing a protein to a customer includes providing a
- 30 pCWin2-MBP-SBD (pMS₃₉) vector as set forth in SEQ ID NO:10 to a protein production facility, wherein a nucleic acid encoding a protein is cloned into the expression vector and the protein is expressed therefrom in the protein production facility, and the protein is provided to a customer. In still another embodiment of the invention, a method of providing a protein to a customer includes providing a

pCWin2-MBP-MCS-SBD (pMXS₃₉) vector as set forth in SEQ ID NO:11 to a protein production facility, wherein a nucleic acid encoding a protein is cloned into the expression vector and the protein is expressed therefrom in the protein production facility, and the protein is provided to a customer.

- 5 In one aspect of the invention, a protein production facility is in-house. In another aspect of the invention, the protein production facility is offsite.

 The invention also includes a method of providing a protein to a customer, wherein at least one glycosyl moiety is added to a protein prior to providing the protein to a customer. In one aspect, a glycosyl moiety is added to a protein in
10 vitro.

 The present invention includes a method of providing a protein to a customer, comprising cloning a nucleic acid encoding said protein into a pCWin1 expression vector as set forth in SEQ ID NO:1, inserting the vector into a bacterial host cell, expressing the protein in the host cell, and providing the protein to a
15 customer.

 Another embodiment of the invention includes a method of providing a protein to a customer, comprising cloning a nucleic acid encoding said protein into a pCWin2 expression vector as set forth in SEQ ID NO:2, inserting the vector into a bacterial host cell, expressing the protein in the host cell, and providing the protein to
20 a customer.

 Another embodiment of the invention includes a method of providing a protein to a customer, comprising cloning a nucleic acid encoding said protein into a pCWin2-MBP expression vector as set forth in SEQ ID NO:3, inserting the vector into a bacterial host cell, expressing the protein in the host cell, and providing the
25 protein to a customer. Yet another embodiment of the invention includes a method of providing a protein to a customer, comprising cloning a nucleic acid encoding said protein into a pCWin2-MBP-SBD (pMS₃₉) expression vector as set forth in SEQ ID NO:10, inserting the vector into a bacterial host cell, expressing the protein in the host cell, and providing the protein to a customer. Still another embodiment of the
30 invention includes a method of providing a protein to a customer, comprising cloning a nucleic acid encoding said protein into a pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector as set forth in SEQ ID NO:11, inserting the vector into a bacterial host cell, expressing the protein in the host cell, and providing the protein to a customer.

In one embodiment of the invention, a method includes adding at least one glycosyl moiety to a protein prior to providing the protein to a customer. In one aspect, a glycosyl moiety is added to a protein in vitro.

The invention features isolated pcWIN1 expression vector comprising
5 the sequence set forth in SEQ ID NO:1. The invention also features an isolated pcWIN1 expression vector consisting of the sequence set forth in SEQ ID NO:1.

In another aspect, the invention features an expression vector comprising the sequence set forth in SEQ ID NO:2. The invention also features an isolated pcWIN2 expression vector consisting of the sequence set forth in SEQ ID
10 NO:2.

In yet another aspect, the invention features an isolated pcWIN2/MBP expression vector comprising the sequence set forth in SEQ ID NO:3. The invention also features an isolated pcWIN2/MBP expression vector consisting of the sequence set forth in SEQ ID NO:3. The invention further features a pcWIN2/MBP expression
15 vector, wherein the pcWIN2/MBP vector comprises a protease cleavage site coding sequence adjacent to the MBP coding sequence.

In another aspect, the invention features an isolated pCWin2-MBP-SBD (pMS₃₉) expression vector comprising the sequence set forth in SEQ ID NO:10. The invention also features an isolated pCWin2-MBP-SBD (pMS₃₉) expression vector
20 consisting of the sequence set forth in SEQ ID NO:10.

In still another aspect, the invention features an isolated pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector comprising the sequence set forth in SEQ ID NO:11. The invention also features an isolated pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector consisting of the sequence set forth in SEQ ID NO:11.
25

The invention features a method of expressing a protein from an isolated pcWIN1 expression vector comprising the sequence set forth in SEQ ID NO:1. In another embodiment, the invention features a method of expressing a protein from an isolated pcWIN2 expression vector comprising the sequence set forth in SEQ ID NO:2. In yet another embodiment, the invention features a method of
30 expressing a protein from an isolated pcWIN2/MBP expression vector comprising the sequence set forth in SEQ ID NO:3. In still another embodiment, the invention features a method of expressing a protein from an isolated pCWin2-MBP-SBD (pMS₃₉) expression vector comprising the sequence set forth in SEQ ID NO:10. In another embodiment, the invention features a method of expressing a protein from an

isolated pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector comprising the sequence set forth in SEQ ID NO:11. In one aspect, the protein is expressed in a prokaryotic cell.

5

BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing summary, as well as the following detailed description of the invention, will be better understood when read in conjunction with the appended drawings. For the purpose of illustrating the invention, there are shown in the drawings embodiment(s) which are presently preferred. It should be understood, however, that invention is not limited to the precise arrangements and instrumentalities shown. In the drawings:

Figure 1A is an image of an electrophoretic gel containing products of a restriction digest. Lanes 1 and 3 are BstEII DNA Marker, lane 2 is SacI/XbaI-digested Cst-04 vector and lane 4 is Kan^r PCR product digested with SacI/XbaI.

Figure 1B is an image of an agar plate showing the result of Cst-04-Kan^r transformation plated on LB kan^r plate.

Figure 1C is an image of an electrophoretic gel containing DNA from an E. coli colony that screened positive for the Cst-04-Kan^r insert. Lane 1 contains BstEII DNA Marker. Lanes 2-4 contain DNA isolated from the Cst-04-Kan5 colony. Lane 2 contains DNA cut with NdeI, lane 3 contains DNA cut with SalI, and lane 4 contains DNA cut with PstI.

Figure 1D is an image of an ampicillin-containing agar plate and a kanamycin-containing agar plate, on both of which Cst-04-Kan5 was streaked. The ampicillin-containing plate inhibited the growth of Cst-04-Kan5-containing cells, demonstrating that the ampicillin gene in Cst-04-Kan5-containing cells is inactive, whereas the kanamycin-containing plate supports the growth of Cst-04-Kan5-containing cells, demonstrating that the kanamycin gene in Cst-04-Kan5-containing cells is operative.

Figure 1E is an image of thin-layer chromatography of the products of the activity of Cst-04Kan5 plasmid-containing cell lysates using lacto-N-neotetraose as a substrate. Lanes labeled 1 and 2 are Cst-04Kan5 from JM109 cells, lanes labeled 3 are Cst-04Kan5 isolated from TG1 cells, and lanes labeled 4 are Cst-04-6-1.

Figure 2A is an image of the agarose gel from which restriction enzyme-digested PCR products were isolated. Lanes marked "M" contain 1 kb DNA

markers, lane 1 contained pCWIN1 insert, lane 2 contained NdeI/ScaI-digested pCWori Kan^r Cst04Kan5 vector, lane 3 contained pre-pCWIN2 insert, and lane 4 contained BamHI/EcoRI-digested pCWori Kan^r Cst04Kan5 vector.

5 Figure 2B is an image of an electrophoretic gel, illustrating the results of restriction digestion of plasmid DNA isolated from positive transformants as a result of pCWin1 and pre-pCWin2 DNA mini-prep. Lanes labeled "M" contain 1 kb DNA markers. Lanes 1 to 5 contain pCWin1 clones. Lanes 6 to 14 contains pre-pCWin2 clones. All 14 clones were digested with EcoRI.

10 Figure 2C is an image of an electrophoretic gel, illustrating the results of restriction digestion of plasmid DNA isolated from positive transformants as a result of pCWin1 and pre-pCWin2 DNA mini-prep. Lanes labeled "M" contain 1 kb DNA markers. Lane 1 contains pCWin1 clone #5, lane 2 contains pre-pCWin2 clone # 11. Both clones # 5 and #11 were digested with NdeI and ScaI.

15 Figure 2D is an image of an electrophoretic gel, illustrating the results of restriction pCWin2 mini-prep screening. Lanes labeled "M" contains 1 kb DNA markers. Lanes 1 through 18 contain pCWin2 clones. The clones were all digested with PstI.

Figure 3A is an image of an electrophoretic gel containing the NdeI/BamHI-digested male cDNA.

20 Figure 3B is an image of an electrophoretic gel containing the restriction enzyme-digested pCWin2 vector.

Figure 3C is an image of two electrophoretic gels containing the restriction enzyme-digested pCWin2 vector. This figure represents the screening of colonies to verify that the male NdeI and BamHI insert size was correct. The first lane on each gel contains 1 kb DNA molecular weight markers, as indicated in the figure. Lanes 1, 2, 3, 4, 5, 7, 8, 9 and 10 correspond to colonies selected from the transformation plate and which positively show the presence of the male cDNA. Lane 6 corresponds to a colony selected from the transformation plate bearing a vector that does not contain the male insert.

30 Figures 4A, 4B, and 4C comprise the entire nucleotide sequence of pCWIN1, as set forth in SEQ ID NO:1.

Figures 5A, 5B, and 5C comprise the entire nucleotide sequence of pCWIN2, as set forth in SEQ ID NO:2.

Figures 6A, 6B, 6C, and 6D comprise the entire nucleotide sequence of pCWIN2-MBP, as set forth in SEQ ID NO:3.

Figure 7A is an image of an electrophoretic gel illustrating the results of a restriction enzyme-digested PCR reaction used to create the SBD₃₉ insert. Lane M is a 1kb DNA marker. Lane 1 is the SBD₃₉ PCR insert product digested with Bgl II and BamHI. The expected size for the SBD₃₉ insert is 447bp.

Figure 7B is an image of an electrophoretic gel illustrating the result of the restriction enzyme digestion of pCWin2-MBP-ST3Gal III (Δ 73) vector. Lane M is a 1kb DNA marker. Lane 1 is pCWin2-MBP-ST3Gal III (Δ 73) digested with BamHI. The expected size for the vector is 7 kb.

Figure 7C is an image of an electrophoretic gel illustrating the results of the DNA mini-prep enzymatic digestion screen of pCWin2-MBP-SBD-ST3Gal III (Δ 73). Lanes M are a 1kb DNA marker, lanes 1 through 11 are pCWin2MBP-SBD-ST3Gal III Δ 73 construct colonies 1 through 11 respectively, digested with Nde I and BamHI. The expected size for the pCWin2-ST3Gal III Δ 73 vector band is 5.9 kb and the expected size for the MBP-SBD insert is 1.6 kb. Clone #6 (Lane 6) illustrates a positive result.

Figure 7D is an image of an electrophoretic gel illustrating the restriction enzyme digestion of pCWin2 vector with BamHI and ScaI. Lane M is a 1kb DNA marker. Lane 1 is digested pCWin2. The expected size for the vector is 4.3 kb and the expected size for the BamH I /Sca I MCS insert is 0.8 kb.

Figure 7E is an image of an electrophoretic gel illustrating the results of the restriction enzyme digestion of pCWin-MBP-SBD₃₉-ST3Gal III (Δ 73). Lane M is a 1kb DNA marker. Lane 1 is pCWin-MBP-SBD₃₉-ST3Gal III (Δ 73) digested with BamH I and Sca I. The expected size for the vector is Linear is pCWin-MBP-SBD₃₉ is 5.8 kb and the expected size for the BamH I /Sca I ST3Gal III (Δ 73) insert is 1.6 kb.

Figure 7F is an image of an electrophoretic gel illustrating the results of the DNA mini-prep restriction enzyme digestion screen of pCWin2-MBP-SBD₃₉ clones. Lanes M is a 1kb DNA marker, lanes 1 through 16 are pCWin2-MBP-SBD₃₉ construct colonies 1 through 16 respectively, digested with Nde I and Xba I. The expected size for the pCWin2 vector band is 5.0 kb and the expected size for the MBP-SBD insert is 1.65 kb. Clone #1 (Lane 1) illustrates a positive result.

Figure 7G is a nucleic acid vector feature map, illustrating restriction sites for the pCWin-MBP-SBD₃₉ (pMS₃₉) construct.

Figure 8A is an electrophoretic gel illustrating the results of the PCR reaction to prepare the SBD insert. Lane M is λ BstE II DNA marker, lane 1 is SBD
5 PCR insert product. The expected size for the SBD insert is 447bp.

Figure 8B is an image of an electrophoretic gel illustrating the results of DNA isolated from PCR-Blunt-SBD colonies and subjected to restriction enzyme digestion. Lane M is a λ BstE II DNA marker, lanes 1 through 8 are PCR-Blunt-SBD colonies 13, 14, 15, 17, 18, 19, 20, and 22 respectively, all digested with XhoI and
10 SalI. The expected size for the PCR-Blunt vector band is 3kb and the expected size for the SBD insert is 447bp.

Figure 8C is an image of an electrophoretic gel illustrating the results of the restriction enzyme-digested pCWin2-MBP kan^r vector. Lane M is a 1kb DNA ladder, lane 1 is pCWin2-MBP kan^r Vector digested with XhoI and Sal I. The
15 expected size for the pCWin2-MBP kan^r vector band is 3kb.

Figure 8D is an image of an electrophoretic gel illustrating the results of DNA isolated from pCWin2-MBP-MCS-SBD (pMXS₃₉) vector-containing colonies. Lane M is a λ BstE II DNA marker, lanes 1 through 13 are pMXS₃₉ colonies 1 through 13 respectively, digested with XhoI and SalI. The expected size
20 for the pCWin2-MBP vector band is 6.1kb and the expected size for the SBD insert is 447bp. Two out of thirteen colonies had the correct size of insert and pMXS₃₉ vector (see lanes 4 and 5 in Figure 8D).

Figure 8E is a nucleic acid vector feature map, illustrating restriction sites for the pCWin-MBP-MCS-SBD₃₉ (pMXS₃₉) construct.

25

DETAILED DESCRIPTION OF THE INVENTION

The use of therapeutic proteins to treat patients experiencing disease or illness increases yearly. Protein therapeutics typically lack the same problematic side effects found with certain traditional chemical therapeutics. Even in instances where
30 the protein therapeutic is altered slightly from its natural state, such a protein typically does not have the same side effects as do certain chemical therapeutics. Similar to the increase in the use of therapeutic proteins, the use of non-therapeutic, or "reagent" proteins increases exponentially from year to year. For example, reagent proteins are

used in such areas as food biochemistry, bioremediation, production of small molecule therapeutics, and even in the production of therapeutic proteins.

The increasing use of protein reagents and therapeutics has enhanced the need for production and preparation of such proteins. It is generally impractical, in terms of cost and time, to isolate and purify a protein therapeutic from its natural source. The cost of isolating proteins from natural sources is prohibitive, and the amount of time needed for such isolation techniques is lengthy. For example, a difficult and time-consuming process for isolating a therapeutic protein from a natural source will drive up the cost of that reagent or therapeutic, which in the latter instance, may unduly burden a medical patient, the patient's insurer, or both. Further, a burdensome isolation process can limit the amount of therapeutic protein available to those in need thereof. Finally, a difficult isolation process can also overburden the entity that produces the reagent or therapeutic protein, reducing profits and wasting valuable business time.

In vitro systems have therefore been developed to produce recombinant forms of reagent proteins and therapeutic proteins. One of the most significant groups of organisms used as an in vitro system for production of recombinant therapeutic proteins is bacteria, and in particular, *Escherichia coli*. *E. coli* is often used for its simplicity, as it is easy to culture and to maintain, and more importantly, it is easy to manipulate genetically. Further, it is relatively simple to isolate protein expressed from *E. coli*.

There are numerous expression vectors that are compatible with bacteria, and in particular, with *E. coli*, for the purpose of producing recombinant therapeutic proteins. However, many vectors are useful only under particular circumstances, and therefore have drawbacks with respect to their utility for selected protein expression under the specific circumstances that may be required. The present invention sets forth methods of providing a protein to a customer that overcome some of the difficulties associated with commercial protein production.

The present invention therefore features a method of providing a protein to a customer, wherein the protein of interest is expressed in a vector containing a nucleic acid encoding the protein of interest, as well as a multiple cloning site and an antibiotic resistance marker, and further wherein the resulting protein is provided to a customer. Part of the advantage of the present invention is that the expression vectors of the present invention reduce the complexity of

subcloning the cDNA encoding a therapeutic protein. Further, the expression vector of the present invention enables the production of proteins using an antibiotic resistance marker other than the ampicillin antibiotic resistance marker, which is not approved for Good Manufacturing Practice (GMP) protocols required by the Food and Drug Administration.

Part of the advantage of the present invention is due to the flexibility of the expression vector used to express the protein. The flexibility of a vector of the present invention provides that a protein can be produced rapidly and efficiently. A cDNA encoding a protein of interest can be readily subcloned into the expression vector by way of a multiple cloning site. Therefore, a method of the present invention features the use of an expression vector as described above to provide a protein to a customer in a more efficient manner.

Another advantage of the present invention is that expression vectors of the invention offer increased productivity and efficiency of protein expression. That is, the design and use of vectors of the present invention provide increased levels of protein expression and production, leading to increased efficiencies in protein expression over similar vectors known in the art. Such advantages increase the quantity of protein produced, and therefore also serve to lower the cost of protein production and increase profit through sales of protein.

The flexibility and functionality of a vector of the present invention increases the ease, efficiency and reliability of the delivery of a protein to a customer. The use of a method of the present invention to streamline and enhance protein product delivery to a customer not only increases the production and profitability of a business entity using such methods, but it also has the effect of increasing the opportunity for medical patients in need thereof with a therapeutic protein.

The present invention also features vectors for expression of reagent proteins, and methods of providing reagent proteins produced using such vectors to a customer. Vectors of the invention useful for the expression of reagent proteins may be the same vectors used to produce therapeutic proteins in methods of the invention. Additionally, vectors of the invention useful for the expression of reagent proteins may be different vectors than those used to produce therapeutic proteins in methods of the invention.

Vectors of the invention designed for production of reagent proteins are further useful for production of proteins that are themselves useful in the

subsequent production of small chemical therapeutics and for remodeling of non-protein molecules, such as carbohydrates. Examples of such proteins include glycosyltransferases, glycosidases and enzymes used in the production of sugar nucleotides. Further, such proteins are useful to produce and remodel carbohydrate-containing glycoproteins. The production and remodeling of glycoproteins has significant therapeutic impact, as glycoproteins form the basis of a significant number of recombinant therapeutics.

Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described herein.

As used herein, each of the following terms has the meaning associated with it in this section.

The articles "a" and "an" are used herein to refer to one or to more than one (i.e. to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

"Encoding" refers to the inherent property of specific sequences of nucleotides in a polynucleotide, such as a gene, a cDNA, or an mRNA, to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (i.e., rRNA, tRNA and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom.

Thus, a gene encodes a protein if transcription and translation of mRNA corresponding to that gene produces the protein in a cell or other biological system. Both the coding strand, the nucleotide sequence of which is identical to the mRNA sequence and is usually provided in sequence listings, and the non-coding strand, used as the template for transcription of a gene or cDNA, can be referred to as encoding the protein or other product of that gene or cDNA.

A "coding region" of a gene consists of the nucleotide residues of the coding strand of the gene and the nucleotides of the non-coding strand of the gene which are homologous with or complementary to, respectively, the coding region of an mRNA molecule which is produced by transcription of the gene.

A "coding region" of an mRNA molecule also consists of the nucleotide residues of the mRNA molecule which are matched with an anticodon region of a transfer RNA molecule during translation of the mRNA molecule or which encode a stop codon. The coding region may thus include nucleotide residues corresponding to amino acid residues which are not present in the mature protein encoded by the mRNA molecule (*e.g.*, amino acid residues in a protein export signal sequence).

An "affinity tag" is a peptide or polypeptide that may be genetically or chemically fused to a second polypeptide for the purposes of purification, isolation, targeting, trafficking, or identification of the second polypeptide. The "genetic" attachment of an affinity tag to a second protein may be effected by cloning a nucleic acid encoding the affinity tag adjacent to a nucleic acid encoding a second protein in a nucleic acid vector.

As used herein, the term "glycosyltransferase," refers to any enzyme/protein that has the ability to transfer a donor sugar to an acceptor moiety.

A "sugar nucleotide-generating enzyme" is an enzyme that has the ability to produce a sugar nucleotide. Sugar nucleotides are known in the art, and include, but are not limited to, such moieties as UDP-Gal, UDP-GlcNAc, and CMP-NAN.

An "isolated nucleic acid" refers to a nucleic acid segment or fragment which has been separated from sequences which flank it in a naturally occurring state, *e.g.*, a DNA fragment which has been removed from the sequences which are normally adjacent to the fragment, *e.g.*, the sequences adjacent to the fragment in a genome in which it naturally occurs. The term also applies to nucleic acids which have been substantially purified from other components which naturally accompany the nucleic acid, *e.g.*, RNA or DNA or proteins, which naturally accompany it in the cell. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule (*e.g.*, as a cDNA or a genomic or cDNA fragment produced by PCR or restriction enzyme digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequence.

In the context of the present invention, the following abbreviations for the commonly occurring nucleic acid bases are used. "A" refers to adenosine, "C" refers to cytidine, "G" refers to guanosine, "T" refers to thymidine, and "U" refers to uridine.

- 5 A "polynucleotide" means a single strand or parallel and anti-parallel strands of a nucleic acid. Thus, a polynucleotide may be either a single-stranded or a double-stranded nucleic acid.

The term "nucleic acid" typically refers to large polynucleotides.

- 10 The term "oligonucleotide" typically refers to short polynucleotides, generally no greater than about 50 nucleotides. It will be understood that when a nucleotide sequence is represented by a DNA sequence (i.e., A, T, G, C), this also includes an RNA sequence (i.e., A, U, G, C) in which "U" replaces "T."

- 15 Conventional notation is used herein to describe polynucleotide sequences: the left-hand end of a single-stranded polynucleotide sequence is the 5' end; the left-hand direction of a double-stranded polynucleotide sequence is referred to as the 5'-direction.

- A first defined nucleic acid sequence is said to be "immediately adjacent to" a second defined nucleic acid sequence when, for example, the last nucleotide of the first nucleic acid sequence is chemically bonded to the first
20 nucleotide of the second nucleic acid sequence through a phosphodiester bond. Conversely, a first defined nucleic acid sequence is also said to be "immediately adjacent to" a second defined nucleic acid sequence when, for example, the first nucleotide of the first nucleic acid sequence is chemically bonded to the last nucleotide of the second nucleic acid sequence through a phosphodiester bond.

- 25 A first defined polypeptide sequence is said to be "immediately adjacent to" a second defined polypeptide sequence when, for example, the last amino acid of the first polypeptide sequence is chemically bonded to the first amino acid of the second polypeptide sequence through a peptide bond. Conversely, a first defined polypeptide sequence is said to be "immediately adjacent to" a second defined
30 polypeptide sequence when, for example, the first amino acid of the first polypeptide sequence is chemically bonded to the last amino acid of the second polypeptide sequence through a peptide bond.

The direction of 5' to 3' addition of nucleotides to nascent RNA transcripts is referred to as the transcription direction. The DNA strand having the

same sequence as an mRNA is referred to as the "coding strand"; sequences on the DNA strand which are located 5' to a reference point on the DNA are referred to as "upstream sequences"; sequences on the DNA strand which are 3' to a reference point on the DNA are referred to as "downstream sequences."

- 5 Unless otherwise specified, a "nucleotide sequence encoding an amino acid sequence" includes all nucleotide sequences that are degenerate versions of each other and that encode the same amino acid sequence. Nucleotide sequences that encode proteins and RNA may include introns.

- "Homologous" as used herein, refers to nucleotide sequence similarity
10 between two regions of the same nucleic acid strand or between regions of two different nucleic acid strands. When a nucleotide residue position in both regions is occupied by the same nucleotide residue, then the regions are homologous at that position. A first region is homologous to a second region if at least one nucleotide residue position of each region is occupied by the same residue. Homology between
15 two regions is expressed in terms of the proportion of nucleotide residue positions of the two regions that are occupied by the same nucleotide residue. By way of example, a region having the nucleotide sequence 5'-ATTGCC-3' and a region having the nucleotide sequence 5'-TATGGC-3' share 50% homology. Preferably, the first region comprises a first portion and the second region comprises a second portion,
20 whereby, at least about 50%, and preferably at least about 75%, at least about 90%, or at least about 95% of the nucleotide residue positions of each of the portions are occupied by the same nucleotide residue. More preferably, all nucleotide residue positions of each of the portions are occupied by the same nucleotide residue.

- As used herein, "homology" is used synonymously with "identity."
25 The determination of percent identity between two nucleotide or amino acid sequences can be accomplished using a mathematical algorithm. For example, a mathematical algorithm useful for comparing two sequences is the algorithm of Karlin and Altschul (1990, Proc. Natl. Acad. Sci. USA 87:2264-2268), modified as in Karlin and Altschul (1993, Proc. Natl. Acad. Sci. USA 90:5873-5877). This
30 algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al. (1990, J. Mol. Biol. 215:403-410), and can be accessed, for example, at the BLAST site of the National Center for Biotechnology Information (NCBI) world wide web site at the National Library of Medicine (NLM) at the National Institutes of Health (NIH). BLAST nucleotide searches can be performed with the NBLAST program

(designated "blastn" at the NCBI web site), using the following parameters: gap penalty = 5; gap extension penalty = 2; mismatch penalty = 3; match reward = 1; expectation value 10.0; and word size = 11 to obtain nucleotide sequences homologous to a nucleic acid described herein. BLAST protein searches can be performed with the XBLAST program (designated "blastn" at the NCBI web site) or the NCBI "blastp" program, using the following parameters: expectation value 10.0, BLOSUM62 scoring matrix to obtain amino acid sequences homologous to a protein molecule described herein.

To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997, Nucleic Acids Res. 25:3389-3402). Alternatively, PSI-Blast or PHI-Blast can be used to perform an iterated search which detects distant relationships between molecules (*id.*) and relationships between molecules which share a common pattern. When utilizing BLAST, Gapped BLAST, PSI-Blast, and PHI-Blast programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used as available on the website of the National Center for Biotechnology Information of the National Library of Medicine at the National Institutes of Health.

The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, typically exact matches are counted.

"Polypeptide" refers to a polymer composed of amino acid residues, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof linked via peptide bonds, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof. Synthetic polypeptides can be synthesized, for example, using an automated polypeptide synthesizer.

As used herein, amino acids are represented by the full name thereof, by the three letter code corresponding thereto, or by the one-letter code corresponding thereto, as indicated in the following table:

	<u>Full Name</u>	<u>Three-Letter Code</u>	<u>One-Letter Code</u>
	Aspartic Acid	Asp	D
	Glutamic Acid	Glu	E
	Lysine	Lys	K
	Arginine	Arg	R

	Histidine	His	H
	Tyrosine	Tyr	Y
	Cysteine	Cys	C
	Asparagine	Asn	N
5	Glutamine	Gln	Q
	Serine	Ser	S
	Threonine	Thr	T
	Glycine	Gly	G
	Alanine	Ala	A
10	Valine	Val	V
	Leucine	Leu	L
	Isoleucine	Ile	I
	Methionine	Met	M
	Proline	Pro	P
15	Phenylalanine	Phe	F
	Tryptophan	Trp	W

The term "protein" typically refers to large polypeptides.

The term "peptide" typically refers to short polypeptides.

20 Conventional notation is used herein to portray polypeptide sequences: the left-hand end of a polypeptide sequence is the amino-terminus; the right-hand end of a polypeptide sequence is the carboxyl-terminus.

A "therapeutic protein" as the term is used herein refers to any protein that is useful to treat a disease state or to improve the overall health of a living
25 organism. A therapeutic protein may effect such changes in a living organism when administered alone, or when used to improve the therapeutic capacity of another substance.

A "reagent protein" as the term is used herein refers to any protein that is useful in food biochemistry, bioremediation, production of small molecule
30 therapeutics, and even in the production of therapeutic proteins. Typically, reagent proteins are enzymes capable of catalyzing a reaction to produce a product useful in any of the aforementioned areas.

A "vector" is a composition of matter which comprises an isolated nucleic acid and which can be used to deliver the isolated nucleic acid to the interior

of a cell. Numerous vectors are known in the art including, but not limited to, linear polynucleotides, polynucleotides associated with ionic or amphiphilic compounds, plasmids, and viruses. Thus, the term "vector" includes an autonomously replicating plasmid or a virus. The term should also be construed to include non-plasmid and non-viral compounds which facilitate transfer of nucleic acid into cells, such as, for example, polylysine compounds, liposomes, and the like. Examples of viral vectors include, but are not limited to, adenoviral vectors, adeno-associated virus vectors, retroviral vectors, and the like.

"Expression vector" refers to a vector comprising a recombinant polynucleotide comprising expression control sequences operatively linked to a nucleotide sequence to be expressed. An expression vector comprises sufficient cis-acting elements for expression; other elements for expression can be supplied by the host cell or in an in vitro expression system. Expression vectors include all those known in the art, such as cosmids, plasmids (e.g., naked or contained in liposomes) and viruses that incorporate the recombinant polynucleotide.

A "multiple cloning site" as the term is used herein is a region of a nucleic acid vector that contains more than one sequence of nucleotides that is recognized by at least one restriction enzyme.

An "antibiotic resistance marker" as the term is used herein refers to a sequence of nucleotides that encodes a protein which, when expressed in a living cell, confers to that cell the ability to live and grow in the presence of an antibiotic.

The term "saccharide" refers in general to any carbohydrate, a chemical entity with the most basic structure of $(CH_2O)_n$. Saccharides vary in complexity, and may also include nucleic acid, amino acid, or virtually any other chemical moiety existing in biological systems.

"Monosaccharide" refers to a single unit of carbohydrate of a defined identity.

"Oligosaccharide" refers to a molecule consisting of several units of carbohydrates of defined identity. Typically, saccharide sequences between 2-20 units may be referred to as oligosaccharides.

"Polysaccharide" refers to a molecule consisting of many units of carbohydrates of defined identity. However, any saccharide of two or more units may correctly be considered a polysaccharide.

A "party" as the term is used herein refers to an individual or an entity involved in a transaction related to a method, a vector, or a protein of the present invention. For example, an individual who provides a vector to a business entity is considered to be a "party" in the context of the present invention. Further, the
5 business entity is also considered to be a "party" in the context of the present invention.

A "recipient" as the term is used herein refers to a specific party who receives a vector or a protein of the present invention. For example, if an individual gives a business entity a vector of the invention, the business entity is considered to be
10 a "recipient" in the context of the present invention. By way of another example, an individual within an organization may provide a second individual within the same organization with a vector or a protein of the present invention. The second individual, who is in receipt of a vector or a protein of the invention, is considered to be a "recipient" in terms of the present invention. It should be noted that a recipient
15 may be a customer, but that not all customers are recipients.

A "customer," as the term is used herein, refers to an intended recipient of a specific item in a formal transaction. A customer is also a recipient, but is distinct from a recipient in that a customer is recipient who is an endpoint for a transaction, whereas a recipient may be an intermediate in a larger transaction. For
20 example, customers of the present invention include, but are not limited to, an entity responsible for the creation of an expression vector that contains the cDNA encoding a protein of interest, if that entity will use the protein produced. A customer is also an entity responsible for expression of a protein from a vector that contains the cDNA encoding a protein of interest, if that entity will use the protein produced. A customer
25 also may be an entity that purchases a protein expressed from a vector of the present invention, for the purpose of using the protein. Further, the entity that creates an expression vector of the present invention may be a customer if that entity uses the protein produced by the vector.

A "protein production facility" as the term is used herein is any
30 location that has the ability to express a protein encoded within a nucleic acid vector.

As the term is used herein, "in-house" refers to dealings within a single organization. In this context, an organization may be a single company, two or more jointly cooperating laboratories, or a corporation and its subsidiaries, collectively.

"Offsite," as the term is used herein, refers to dealings that extend beyond the "in-house" context discussed above. For example, the transfer of a vector of the invention from a first organization to a second organization is a transfer of that vector "offsite."

5

I. Vectors

The present invention includes an isolated nucleic acid encoding a protein operably linked to a nucleic acid comprising a promoter/regulatory sequence such that the nucleic acid is capable of directing expression of the protein encoded by the nucleic acid. Thus, the invention encompasses expression vectors and methods for the expression of proteins based on exogenous DNA introduced into cells with concomitant expression of the exogenous DNA in the cells such as those described, for example, in Sambrook et al. (1989, *supra*), and Ausubel et al. (1997, *supra*).

15 An expression vector of the present invention is based on the pcWori+ vector (Muchmore et al., 1987, *Meth. Enzymol.* 177:44-73). However, the pcWori+ vector by itself is not adequate for the production of protein reagents and therapeutic proteins according to the present invention. The pcWori+ vector contains an ampicillin resistance marker. Certain regulatory agencies require that the production of proteins for therapeutic use cannot be carried out using recombinant vectors containing ampicillin resistance genes. Therefore, an expression vector of the present invention features an antibiotic resistance marker approved by the U.S. Food and Drug Administration (FDA) for use in the production of protein reagents and therapeutic proteins. Such antibiotic resistance markers include, but are not limited to, kanamycin, tetracycline, and chloramphenicol.

25 In the invention, the ampicillin resistance marker normally present in the pcWori+ vector is disabled as follows. Briefly, the ampicillin resistance marker in the pcWori+ vector is disrupted in order to produce, in part, a vector of the present invention. PCR primers designed to create PvuI and ScaI restriction enzyme cleavage sites on either end of a kanamycin resistance gene are used, and the resultant PCR product is digested with PvuI and ScaI restriction enzymes. The ampicillin resistance gene in a pcWori+ vector is also cut with PvuI and ScaI restriction enzymes. Subsequently, the kanamycin resistance gene is ligated into the pcWori+ vector that was cleaved within the ampicillin resistance gene.

30

Verification of successful disruption of the ampicillin resistance gene and successful insertion of the kanamycin resistance gene is observed by transforming *E. coli* cells, for example, with the ligation mixture. Growth of the transformed cells on kanamycin-containing agar plates confirms the successful insertion of the

5 kanamycin resistance marker, while lack of growth on ampicillin containing plates confirms the successful disruption of the ampicillin resistance gene. Other methods of disruption or deletion of the ampicillin resistance gene will be known to one of skill in the art. Similarly, other methods of inserting the kanamycin resistance gene, or any other antibiotic resistance gene useful in the present invention, and methods of

10 confirming the insertion and/or deletion of genes will also be known to one of skill in the art.

Another feature of a vector of the present invention is a versatile and highly-functional multiple cloning site. As described in, for example, in Sambrook et al. (1989, *supra*), a "multiple cloning site" is a nucleic acid having a sequence

15 encoding more than one restriction enzyme recognition site. The practical purpose of a multiple cloning site is to allow the ligation (i.e., "insertion") of an exogenous polynucleotide into the multiple cloning site, wherein the exogenous polynucleotide may have different restriction enzyme recognition sequences at its 5' and 3' ends. That is, the multiple cloning site allows flexibility with respect to the identity of the 5'

20 and 3' ends on an exogenous polynucleotide, thus facilitating the cloning of such a polynucleotide into the multiple cloning site.

A multiple cloning site is most often found, and is most useful, in a nucleic acid vector such as a vector of the present invention. As will be known to the skilled artisan, a multiple cloning site may be located adjacent to other functional

25 elements in a vector, such as a promoter. A multiple cloning site may also be designed such that insertion of an exogenous polynucleotide into the multiple cloning site results in the exogenous polynucleotide being expressed in frame with the adjacent elements to create a fusion protein of the protein encoded by the exogenous polynucleotide and the protein encoded by the adjacent element.

30 Accordingly, a vector of the present invention contains at least one multiple cloning site. The creation of a functional multiple cloning site in a vector of the present invention is described in greater detail elsewhere herein. Briefly, a multiple cloning site may be designed and synthesized *de novo*, or it may be isolated from another pre-existing vector. PCR methods are used to create multiple cloning

site polynucleotides having specific restriction enzyme recognition sites on either end (5' and 3') of the multiple cloning site polynucleotide. A multiple cloning site polynucleotide is then inserted into a pcWori+ vector of the present invention by means of specific restriction enzyme recognition sites corresponding to those on either end of the multiple cloning site polynucleotide. As will be understood by one skilled in the art, various molecular biological techniques are available to insert, delete, and/or modify a multiple cloning site in a vector of the present invention in order to create a more functional and flexible multiple cloning site useful in connection with the present invention.

Another feature of a vector of the present invention is the option of an affinity tag coding sequence located in the multiple cloning site. An affinity tag coding sequence may be inserted into the multiple cloning site adjacent to, upstream from, or downstream from a target protein coding sequence. As will be understood by one of skill in the art, an affinity tag will typically be inserted into the multiple cloning site in frame with the target protein. One of skill in the art will also understand that an affinity tag coding sequence can be used to produce a recombinant fusion protein by concomitantly expressing the affinity tag and target protein. The expressed fusion protein can then be isolated, purified, or identified by means of the affinity tag. An affinity tag is especially important when expressing proteins that are reagents and less important when expressing therapeutic proteins due to restrictions imposed by regulatory agencies.

Affinity tags useful in the present invention include, but are not limited to, a maltose binding protein, a histidine tag, a Factor IX tag, a glutathione-S-transferase tag, a FLAG-tag, and a starch binding domain tag. Other tags are well known in the art, and the use of such tags in the present invention would be readily understood by the skilled artisan.

Any single vector of the present invention may have more than one feature described herein. By way of a non-limiting example, a vector of the present invention may have a disrupted ampicillin resistance gene, a functional kanamycin resistance gene, and a modified, multi-functional multiple cloning site. An example of one such vector of the present invention is pCWin1, the sequence of which is set forth in SEQ ID NO:1. A pCWin1 vector of the present invention has, for example, two BamHI restriction enzyme recognition sites, one of which is located within the multiple cloning site. Another example of a vector of the present invention is

pCWin2, the sequence of which is set forth in SEQ ID NO:2. A pCWin2 vector of the present invention has, for example, only one BamHI restriction enzyme recognition site which is located within the multiple cloning site.

- A further example of a vector of the present invention is pCWin2-MBP, the sequence of which is set forth in SEQ ID NO:3. A pCWin2-MBP vector of the invention has, for example, one BamHI restriction enzyme recognition site located within the multiple cloning site and additionally has an E. coli malE maltose binding protein coding sequence inserted into the multiple cloning site in between the NdeI and BamHI restriction enzyme recognition sites. The NdeI sequence in the multiple cloning site contains an ATG start codon. The pCWin2-MBP vector is therefore useful, for example, for expression of a fusion protein comprised of a maltose binding protein and a desired protein. This is achieved by inserting a polynucleotide encoding the desired protein into the multiple cloning site in frame with the maltose binding protein and expressing the entire open reading frame encoded in the multiple cloning site.

- Yet another example of a vector of the present invention is pCWin2-MBP-SBD₃₉ (pMS₃₉), the sequence of which is set forth in SEQ ID NO:10. A pCWin2-MBP-SBD₃₉ (pMS₃₉) vector of the invention has, for example, one BamHI restriction enzyme recognition site located within the multiple cloning site, and one EcoRI restriction enzyme recognition site located within the multiple cloning site, and additionally has an E. coli malE maltose binding protein coding sequence inserted into the multiple cloning site in between the NdeI and SacI restriction enzyme recognition sites. The pCWin2-MBP-SBD₃₉ (pMS₃₉) vector also has a starch-binding domain (SBD) inserted between the EcoRI and BamHI restriction sites. The NdeI sequence in the multiple cloning site contains an ATG start codon. The pCWin2-MBP-SBD₃₉ (pMS₃₉) vector is therefore useful, for example, for expression of a fusion protein comprised of a maltose binding protein, a starch binding domain, and a desired protein. This is achieved by inserting a polynucleotide encoding the desired protein into the multiple cloning site in frame with the maltose binding protein and starch binding domain, and expressing the entire open reading frame encoded in the multiple cloning site.

Still another example of a vector of the present invention is pCWin2-MBP-MCS-SBD₃₉ (pMXS₃₉), the sequence of which is set forth in SEQ ID NO:11. As compared to, the pMXS₃₉ vector expresses, in one aspect of the invention, a fusion

protein with the structure "MBP—desired protein—SBD," whereas the pMS₃₉ vector expresses, in another aspect of the invention, a "MBP—SBD—desired protein" fusion protein. Accordingly, a pCWin2-MBP-SBD₃₉ (pMXS₃₉) vector of the invention has, for example, one XhoI restriction enzyme recognition site located within the multiple cloning site, and one SalI restriction enzyme recognition site located within the multiple cloning site, and additionally has an E. coli malE maltose binding protein coding sequence inserted into the multiple cloning site in between the NdeI and SacI restriction enzyme recognition sites. The pCWin2-MBP-SBD₃₉ (pMXS₃₉) vector also has a starch-binding domain (SBD) inserted between the XhoI and SalI restriction sites. The NdeI sequence in the multiple cloning site contains an ATG start codon. The pCWin2-MBP-SBD₃₉ (pMXS₃₉) vector is therefore useful, for example, for expression of a fusion protein comprised of a maltose binding protein, a starch binding domain, and a desired protein. This is achieved by inserting a polynucleotide encoding the desired protein into the multiple cloning site in frame with the maltose binding protein and starch binding domain, and expressing the entire open reading frame encoded in the multiple cloning site.

A vector of the present invention, as described above, is useful for the production of a therapeutic protein. A polynucleotide sequence encoding a therapeutic protein may be inserted into the multiple cloning site using any technique known to the skilled artisan. For example, a polynucleotide sequence encoding a therapeutic protein may be modified to contain specific restriction enzyme recognition sites at the 5' and 3' ends of the polynucleotide. Such restriction enzyme recognition sites will correspond to recognition sites located within the multiple cloning site of a vector of the present invention, facilitating the insertion (by ligation) of the therapeutic protein-encoding sequence into the multiple cloning site, and when expressed, producing a therapeutic protein.

Therapeutic proteins useful in the present invention are numerous and are well-known in the art, and are therefore not listed here. By way of a non-limiting example, such therapeutic proteins include erythropoietin, human growth hormone, granulocyte colony stimulating factor, interferons alpha, -beta, and -gamma, Factor IX, follicle stimulating hormone, interleukin-2, erythropoietin, anti-TNF-alpha, and a lysosomal hydrolase. Lysosomal hydrolases useful in the present invention include, but are not limited to, beta-glucosidase, alpha-galactosidase-A, beta-hexosaminidase, beta-galactosidase, alpha-galactosidase, alpha-mannosidase, beta-mannosidase, alpha-

L-fucosidase, beta-glucuronidase, alpha-glucosidase, alpha-N-acetylgalactosaminidase, and acid phosphatase. It will be understood that any mutant or variant of a therapeutic protein may be expressed using vectors of the present invention.

5 The present invention also features a vector useful for the production of a non-therapeutic protein, referred to herein as reagent proteins. As will be understood by the skilled artisan, a reagent protein is one which does not currently have a therapeutic application. Such proteins include, but are not limited to, enzyme reagents, food enzymes, nutritional supplements, and non-active additives. Methods
10 of expressing reagent proteins using vectors of the invention will be understood by the skilled artisan to be conducted in the same manner as the above-described methods of expressing therapeutic proteins using vectors of the present invention.

 Another feature of a vector of the present invention is the option of a protease cleavage site coding sequence located in the multiple cloning site. A
15 protease cleavage site coding sequence may be inserted into the multiple cloning site adjacent to, upstream from, or downstream from a target protein coding sequence. As will be understood by one of skill in the art, a protease cleavage site will typically be inserted into the multiple cloning site in frame with the target protein. One of skill in the art will also understand that a protease cleavage site coding sequence can be used
20 to produce a recombinant fusion protein by concomitantly expressing the protease cleavage site sequence and target protein. The expressed fusion protein can then be isolated, purified, or identified by means of the protease cleavage site.

 In an embodiment of the invention, a vector contains a coding sequence for a protease cleavage site which is located C-terminal to the nucleic acid
25 sequence encoding an MBP. In one aspect, a vector is pCWin2-MBP-SBD₃₉ (pMS₃₉). In another aspect, a vector is pCWin2-MBP-MCS-SBD₃₉ (pMXS₃₉).

 A fusion protein containing a preselected protease cleavage site, as will be understood by one of skill in the art, is useful for the removal of amino acid sequence that is extraneous or non-essential to the expressed protein of interest. For
30 example, a target protein may be expressed using a vector of the present invention as a fusion with an affinity tag for the purpose of purification of the target protein, but the affinity tag may not be desirable once the protein is sufficiently purified. The insertion of a specific protease cleavage site between the target protein and the affinity tag is useful for the cleavage of the affinity tag from the target protein.

Protease cleavage sites useful in the present invention include, but are not limited to, an enterokinase cleavage site, a Factor Xa cleavage site, a thrombin cleavage site, and a TEV protease cleavage site. The skilled artisan will understand the characteristics and uses of a protease cleavage useful in the present invention.

5 The present invention also features a recombinant bacterial host cell comprising, *inter alia*, a nucleic acid vector as described elsewhere herein. In one aspect, the recombinant cell is transformed with a vector of the present invention. The transformed vector need not be integrated into the cell genome nor does it need to be expressed in the cell. However, the transformed vector will be capable of being
10 expressed in the cell. In one aspect of the invention, *E. coli* is used for transformation of a vector of the present invention and expression of protein therefrom. In another aspect of the invention, a K-12 strain of *E. coli* is useful for expression of protein from a vector of the present invention. Strains of *E. coli* useful in the present invention include, but are not limited to, JM83, JM101, JM103, JM109, W3110,
15 chi1776, and JA221.

 It will be understood that a host cell useful in the present invention will be capable of growth and culture on a small scale, medium scale, or a large scale. For example, a host cell of the invention is useful for testing the expression of a protein from a vector of the invention equally as much as it is useful for large scale
20 production of a reagent or therapeutic protein product. Techniques useful in culturing host cells and expressing protein from a vector contained therein are well known in the art and will therefore not be listed herein.

 A host cell of the present invention may be transformed with a vector of the present invention to produce a transformed host cell of the invention.
25 Transformation, as known to the skilled artisan, includes the process of inserting a nucleic acid vector into a host cell, such that the host cell containing the nucleic acid vector remains viable. Such transformation of nucleic acid into a bacterial cell is useful for purposes including, but not limited to, creation of a stably-transformed host cell, making a biological deposit, propagating the vector-containing host cell,
30 propagating the vector-containing host cell for the production and isolation of additional vector, expression of target protein encoded by vector, and the like.

 Methods of transforming a vector are numerous and well-known in the art, and will therefore not be listed here. By way of a non-limiting example, a competent bacterial cell of the invention may be transformed by a vector of the

- invention using electroporation. Methods of making bacterial cells "competent" are well-known in the art, and typically involve preparation of the bacterial cells so that the cells take up exogenous DNA. Similarly, methods of electroporation are known in the art, and detailed descriptions of such methods may be found, for example, in
- 5 Sambrook et al. (1989, *supra*). The transformation of a competent cell with vector DNA may be also accomplished using chemical-based methods. One example of a well-known chemical-based method of bacterial transformation is described by Inoue, et al. (1990, *Gene* 96:23-28). Other methods of transformation will be known to the skilled artisan.
- 10 In one embodiment of the present invention, a Cst-04Kan5 plasmid is transformed into *E. coli* JM109 cells using 20 μ l JM109 competent cells in 0.34 μ l 1.42 M beta mercaptoethanol, incubated on ice for 10 minutes, at which time 1 μ l (100 ng) Cst-04-Kan5 plasmid is added to the transformation mixture. The cell/DNA mixture is incubated ice for 30 minutes, then heat shocked at 42 °C for 45 seconds.
- 15 The reaction is then incubated on ice for 2 minutes, at which time 80 μ l SOC media is added. The reaction mixture is then shaken at 37°C for 1 hour, and subsequently, plated on LB Kan^r agar plates. Identification and confirmation of the Cst-04-Kan5 plasmid DNA is carried out using a restriction enzyme digestion of plasmid DNA isolated from positive transformants, using NdeI, SalI, PstI restriction enzymes.
- 20 A transformed host cell of the present invention may be used to express a protein. In an embodiment of the invention, a transformed host cell contains a vector of the invention, which contains therein a nucleic acid sequence encoding an exogenous protein. The protein is expressed using any expression method known in the art (for example, IPTG). The expressed protein may be contained within the host cell, or it may be secreted from the host cell into the growth medium.
- 25 Methods for isolating an expressed protein are well-known in the art, and the skilled artisan will know how to determine the best method for isolation of an expressed protein based on the characteristics of any given host cell expression system. By way of a non-limiting example, an expressed protein that is secreted from a host cell may be isolated from the growth medium. Isolation of a protein from a growth medium may include removal of bacterial cells and cellular debris. By way of another non-limiting example, an expressed protein that is contained within a host cell may be isolated from the host cell. Isolation of such an "intracellular" expressed
- 30

protein may include disruption of the host cell and removal of cellular debris from the resultant mixture. These methods are not intended to be exclusive representations of the present invention, but rather, are merely for the purposes of illustration of various applications of the present invention.

5 Purification of a protein expressed in accordance with the present invention may be effected by any means known in the art. The skilled artisan will know how to determine the best method for the purification of a protein expressed in accordance with the present invention. A purification method will be chosen by the skilled artisan based on factors such as, but not limited to, the expression host, the
10 contents of the crude extract of the protein, the size of the protein, the properties of the protein, the desired end product of the protein purification process, and the subsequent use of the end product of the protein purification process.

 In an embodiment of the invention, isolation or purification of a protein expressed in accordance with the present invention may not be desired. In an
15 aspect of the present invention, an expressed protein may be stored or transported inside the bacterial host cell in which the protein was expressed. In another aspect of the invention, an expressed protein may be used in a crude lysate form, which is produced by lysis of a host cell in which the protein was expressed. In yet another
20 embodiment of the invention, an expressed protein may be partially isolated or partially purified according to any of the methods set forth or described herein. The skilled artisan will know when it is not desirable to isolate or purify a protein of the invention, and will be familiar with the techniques available for the use and preparation of such proteins.

25 II. Methods of providing a protein to a customer

 The present invention features a method of providing a protein to a customer. In an embodiment of the invention, a nucleic acid encoding a protein is cloned into an expression vector. The encoded protein is expressed from the expression vector, and the resulting protein product is provided to a customer.

30 In an embodiment of the invention, a protein is expressed from an expression vector in vitro. Techniques for in vitro protein expression are known in the art, and are exemplified by the methods of Melton and colleagues (Krieg et al., 1987,

Meth. Enzymol. 155, 397-415; Yisraeli et al., 1989, Meth. Enzymol. 180, 42-50). A protein produced using an in vitro expression method of the present invention is provided to a customer.

5 In another embodiment of the invention, a protein is expressed from an expression vector in vivo. Numerous techniques for expression of a protein from an expression vector in vivo are described in detail elsewhere herein, and are also well-known in the art. Such techniques include, but are not limited to, expression of a protein from a vector in a bacterial host cell. A protein produced using an in vivo expression method of the present invention is provided to a customer.

10 In one aspect of the invention, a protein is expressed from a pcWIN1 expression vector, as set forth in SEQ ID NO:1. In another aspect of the invention, a protein is expressed from a pcWIN2 expression vector, as set forth in SEQ ID NO:2. In yet another aspect of the invention, a protein is expressed from a pcWIN2/MBP expression vector, as set forth in SEQ ID NO:3. In still another aspect of the
15 invention, a protein is expressed from a pCWin2-MBP-SBD₃₉ (pMS₃₉) expression vector, as set forth in SEQ ID NO:10. In yet another aspect of the invention, a protein is expressed from a pCWin2-MBP-MCS-SBD₃₉ (pMXS₃₉) expression vector, as set forth in SEQ ID NO:11. As will be understood by one of skill in the art, a pcWIN vector of the present invention is useful in any of the expression methods set forth
20 herein for the production of a target protein that may be provided to a customer.

Methods of the present invention for in vivo expression of a protein in a bacterial cell comprise transformation of the bacterial cell with an expression vector comprising the protein of interest. Methods of transforming a bacterial cell with a vector are described in detail elsewhere herein, and would be understood by one of
25 ordinary skill in the art. It will be appreciated that methods of bacterial cell transformation other than those explicitly disclosed herein are useful in methods of the present invention, and therefore, are within the scope of the present invention.

Vectors featured in methods of the present invention are described in detail elsewhere herein. In an embodiment of the invention, a method of providing a
30 protein to a customer comprises expressing a protein from an expression vector useful for production of a therapeutic protein. As described above, vectors of the invention useful in such methods are comprised of an antibiotic resistance marker such as kanamycin, tetracycline, chloramphenicol, and the like, as such antibiotics are particularly useful in connection with the expression of therapeutic proteins. A

therapeutic protein provided to a customer using this method is particularly useful to the customer due to the fact that kanamycin, tetracycline, chloramphenicol and like antibiotics are preferred by certain regulatory agencies for the production of therapeutic proteins.

5 The present invention therefore features a method of providing a known therapeutic protein to a customer. Therapeutic proteins include, but are not limited to, human growth hormone, granulocyte colony stimulating factor, interferons alpha, -beta, and -gamma, Factor IX, follicle stimulating hormone, beta-glucosidase, interleukin-2, erythropoietin, alpha-galactosidase-A, and anti-TNF-alpha. It will be
10 understood by the skilled artisan that any nucleic acid encoding a therapeutic protein, wherein the nucleic acid is capable of being cloned into and expressed from a nucleic acid vector of the invention, will be useful in the present invention. The ability to determine a therapeutic protein useful in the present invention is within the skill of the ordinary artisan and such a determination does not require undue experimentation.

15 In another embodiment of the invention, a method of providing a protein to a customer comprises expressing a protein from an expression vector useful for production of a reagent protein. Vectors of the invention useful in such methods preferably have had the native ampicillin resistance gene disabled, altered, or deleted, such that the ampicillin resistance gene is no longer functional in the vector. Such
20 vectors are comprised of any antibiotic resistance marker other than ampicillin known in the art to be useful in the expression of proteins. Antibiotic resistance markers useful in the invention include kanamycin, tetracycline, chloramphenicol, and like antibiotic resistance markers approved by certain regulatory agencies, as well as any antibiotic resistance marker not approved by certain regulatory agencies for use with
25 therapeutic proteins.

 Therefore, the present invention also features methods of providing a protein to a customer, wherein the protein is a reagent protein, and therefore, need not be expressed from a vector containing antibiotic resistance marker accepted by a regulatory agency. However, a reagent protein may also be expressed from a vector
30 containing an FDA-accepted antibiotic resistance marker. A protein produced by such a method of the invention may be useful for almost any purpose, including, but not limited to, an enzyme reagent, a food enzyme, a nutritional supplement, and a non-active additive. Examples of such proteins include, but are not limited to a glycosyltransferase and a sugar nucleotide-generating enzyme.

In an embodiment of the invention, a method is provided wherein a nucleic acid is cloned into a vector containing any antibiotic resistance marker useful in the expression of a protein, the protein is expressed therefrom, and the resulting protein product is provided to a customer. It will be understood that such a protein may be expressed in vivo or in vitro.

Methods of the present invention also feature a vector comprising a highly-functional multiple cloning site. Such vectors are described in detail elsewhere herein. In an embodiment of the invention, a method of providing a protein to a customer comprises expressing a protein from an expression vector useful for production of a therapeutic protein. As described above, vectors of the invention useful in such methods are comprised of a highly-functional multiple cloning site, in addition to an antibiotic resistance marker such as kanamycin, tetracycline, chloramphenicol, and the like. In this embodiment, a therapeutic protein is expressed cloned into a vector comprising a highly-functional multiple cloning site in addition to an antibiotic resistance marker, expressed therefrom, and provided to a client. In one aspect of the invention, the multiple cloning site contains at least one of NdeI, BamHI, SacI, HindIII, XbaI, XhoI, EcoRI, KpnI, and SalI restriction enzyme cleavage sites.

Methods of the present invention also feature a vector comprising an affinity tag. Such vectors are described in detail elsewhere herein. In an embodiment of the invention, a method of providing a protein to a customer comprises expressing a protein from an expression vector useful for production of a therapeutic protein. As described above, vectors of the invention useful in such methods are comprised of an affinity tag, in addition to an antibiotic resistance marker such as kanamycin, tetracycline, chloramphenicol, and the like. In this embodiment, a therapeutic protein is expressed cloned into a vector comprising an affinity tag in addition to an antibiotic resistance marker, expressed therefrom, and provided to a client. In a preferred embodiment, the affinity tag is a maltose-binding protein.

In other embodiments of the invention, a useful affinity tag may be, but is not limited to, a histidine tag, a Factor IX tag, a glutathione-S-transferase tag, starch-binding domain, a FLAG-tag, and the like. One of skill in the art will understand that any affinity tag capable of being used with a vector of the present invention will be useful in methods of the invention. Further, the skilled artisan will also appreciate that a single vector of the invention may comprise more than one

affinity tag, and that multiple affinity tags may be identical or may be heterogeneous in sequence.

5 The present invention also features a method of providing a protein to a customer, wherein an expression vector used to express a protein has multiple characteristics as described elsewhere herein. For example, a method of the present invention for providing a protein to a customer comprises the cloning of a nucleic acid encoding the protein into an expression vector, wherein the expression vector comprises a kanamycin resistance marker and a highly-functional multiple cloning site.

10 In an embodiment of the invention, a method of providing a protein to a customer comprises the cloning of a protein into the expression vector set forth in SEQ ID NO:1. The protein-SEQ ID NO:1 construct is transformed into an E. coli cell, the protein is expressed therefrom, and the protein product is provided to the customer. In one aspect of the invention, the E. coli cell is a JM109 cell. In another aspect of the invention, the protein is a therapeutic protein. In yet another aspect of the protein, the protein is a reagent protein.

15 In another embodiment of the invention, a method of providing a protein to a customer comprises the cloning of a protein into the expression vector set forth in SEQ ID NO:2. The protein-SEQ ID NO:2 construct is transformed into an E. coli cell, the protein is expressed therefrom, and the protein product is provided to the customer. In one aspect of the invention, the E. coli cell is a JM109 cell. In another aspect of the invention, the protein is a therapeutic protein. In yet another aspect of the protein, the protein is a reagent protein.

20 In another embodiment of the invention, a method of providing a protein to a customer comprises the cloning of a protein into the expression vector set forth in SEQ ID NO:3. The protein-SEQ ID NO:3 construct is transformed into an E. coli cell, the protein is expressed therefrom, and the protein product is provided to the customer. In one aspect of the invention, the E. coli cell is a JM109 cell. In another aspect of the invention, the protein is a therapeutic protein. In yet another aspect of the protein, the protein is a reagent protein.

25 In still another embodiment of the invention, a method of providing a protein to a customer comprises the cloning of a protein into the expression vector set forth in SEQ ID NO:10. The protein-SEQ ID NO:10 construct is transformed into an E. coli cell, the protein is expressed therefrom, and the protein product is provided to

the customer. In one aspect of the invention, the E. coli cell is a JM109 cell. In another aspect of the invention, the protein is a therapeutic protein. In yet another aspect of the protein, the protein is a reagent protein.

In another embodiment of the invention, a method of providing a protein to a customer comprises the cloning of a protein into the expression vector set forth in SEQ ID NO:11. The protein-SEQ ID NO:11 construct is transformed into an E. coli cell, the protein is expressed therefrom, and the protein product is provided to the customer. In one aspect of the invention, the E. coli cell is a JM109 cell. In another aspect of the invention, the protein is a therapeutic protein. In yet another aspect of the protein, the protein is a reagent protein.

The present invention features a method of providing a protein to a customer, wherein a nucleic acid encoding a protein is cloned into an expression vector of the invention by the party providing a vector to a recipient. That is, a nucleic acid encoding a protein is cloned into an expression vector of the invention before the vector is transferred to a recipient. In an embodiment of the invention, a method of providing a protein to a customer comprises providing a vector to a recipient, wherein the vector contains a nucleic acid encoding a protein. The recipient of the vector expresses the protein, and the protein is then provided to a customer. In one aspect of the invention, a recipient is a protein production facility.

In an embodiment of the invention, a method of providing a protein to a customer comprises providing a vector to a recipient, wherein the vector does not contain a nucleic acid encoding a protein. The recipient of the vector clones a nucleic acid encoding a protein into the vector and expresses the protein, and the protein is then provided to a customer. In one aspect of the invention, a recipient is a protein production facility. In another aspect of the invention, the nucleic acid cloned into a vector is provided by the party providing the vector to the recipient. In yet another aspect of the invention, the nucleic acid cloned into a vector is provided by the recipient.

The invention also features a method of providing a protein to a customer, wherein the method comprises providing a vector to a recipient, wherein the vector comprises a nucleic acid encoding a protein, for the purpose of expression of the protein encoded by the vector. In one embodiment of the invention, the recipient is a protein production facility. In one aspect, the protein production facility is in-house. By way of non-limiting examples, such recipients include an in-house

protein production facility and an in-house laboratory. In another aspect of the invention, the protein production facility is offsite. By way of non-limiting examples, such recipients include an offsite protein production facility, an offsite laboratory, an offsite biotechnology company and an offsite pharmaceutical company.

5 A protein produced using a method of the present invention may be provided to a customer by the recipient of the vector, wherein the recipient is responsible for expressing the protein from the vector provided by another party. Alternatively, a protein produced using a method of the present invention may be provided to a customer by the original party that provided the vector to the recipient,
10 wherein the recipient expresses the protein and provides the resulting protein product to the original party so that the original party may provide the protein to a customer.

 A protein produced using a method of the present invention may be provided to a customer in the form of a purified protein, a partially purified protein, an isolated protein, a partially isolated protein, a bacterial cell lysate, cell paste or
15 purified inclusion bodies. It will be understood that a protein produced using a method of the present invention may be provided to a customer in any form known in the art to be useful for the storage, transfer, or processing of a recombinant protein.

 The present invention also features a method of providing a protein to a customer, wherein at least one glycosyl moiety is added to the protein before
20 providing the protein to the customer. A glycosyl moiety may be added to a protein using any method known in the art. Additionally, a glycosyl moiety may be added to a protein of the invention using any one of the methods or reagents taught by DeFrees et al. in PCT Application WO 03/031464, which is incorporated herein by reference in its entirety. The skilled artisan will understand, based on the disclosure herein, that
25 any of the methods known in the art or set forth herein are useful for glycosylating a protein of the present invention prior to providing the protein to a customer.

 Thus, in an embodiment of the present invention, a method of providing a protein to a customer comprises providing a vector comprising a nucleic acid encoding a protein to a recipient, wherein the recipient expresses the protein, and
30 further wherein the protein is modified with at least one glycosyl moiety before the protein is provided to a customer. In one aspect of the invention, at least one glycosyl moiety is added to the protein by the recipient before providing the protein to a customer. In another aspect of the invention, at least one glycosyl moiety is added to the protein by the recipient before providing the protein to the original supplier of the

vector, wherein the original supplier of the vector subsequently provides the protein to a customer. In yet another aspect of the invention, the recipient provides the expressed protein to the original supplier of the vector, wherein the original supplier of the vector adds at least one glycosyl moiety to the protein before providing the protein to a customer.

In another embodiment of the invention, a method of providing a protein to a customer comprises providing a vector to a recipient, wherein the recipient clones a nucleic acid encoding a protein into the vector, expresses the protein, and further wherein the protein is modified with at least one glycosyl moiety before the protein is provided to a customer. In one aspect of the invention, at least one glycosyl moiety is added to the protein by the recipient before providing the protein to a customer. In another aspect of the invention, at least one glycosyl moiety is added to the protein by the recipient before providing the protein to the original supplier of the vector, wherein the original supplier of the vector to the recipient provides the protein to a customer. In yet another aspect of the invention, the recipient provides the expressed protein to the original supplier of the vector, wherein the original supplier of the vector adds at least one glycosyl moiety to the protein before providing the protein to a customer.

EXPERIMENTAL EXAMPLES

The invention is now described with reference to the following examples. These examples are provided for the purpose of illustration only and the invention should in no way be construed as being limited to these examples but rather should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

Example 1: Modification of pCWori+ Amp^r expression vector by disrupting the Amp^r gene and adding the kanamycin resistance gene

The pCWori+ Amp^r vector contains an ampicillin resistance marker, as well as the genes encoding N. Meningitidis CMP-NAN synthetase (CNS) and Campylobacter Jejuni α 2,3 Sialyl Transferase (CstI), referred to as Cst-04 (Cst-04 was provided by Warren Wakarchuck, National Research Council, Canada). This

example describes the complete process by which the Cst-04 (pCWori+ amp^r-CNS-CstI) plasmid was interrupted at the PvuI and ScaI sites of ampicillin gene by the insertion of the kanamycin resistance gene.

- A kanamycin resistance gene was isolated from pGEX-Kt-ext Kan^r using PCR to generate cDNA with modified restriction sites at 5' (PvuI-ATTCCAATTTCGATCGGGGGGGGGGAAA) (SEQ ID NO:4) and 3' (ScaI-ATTCCAAGTAGTACTTTAGAAAACTCATCG) (SEQ ID NO:5) ends. The PCR product was then subcloned into a Cst04 (pCWori+ amp^r-CNS-CstI) vector in TG1 cells. A colony positive for the recombinant vector (Cst-04Kan5) was identified, and the Cst-04Kan5 plasmid was isolated, then transformed into JM109 cells.

- A PCR reaction was conducted containing 1 ng pGEXKan^r template, 1 µg (1 µl) kan-ScaI/PvuI primer, 77 µl H₂O, 8 µl dNTP mixture, 10 µl 10X buffer, and 1 µl Vent polymerase. The reaction parameters included a 5 minute cycle at 95 °C, followed by the addition of 1 µl of Vent polymerase and thirty cycles of the following temperature pattern: 94 °C for one minute, 55 °C for one minute, 72 °C for one minute.

- The PCR product and the Cst-04 vector were subjected to a restriction digest. The PCR product digest included 16 µl PCR rxn, 2 µl 10X buffer, 0.5 µl PvuI, 0.5 µl ScaI, 1 µl H₂O. The pGEX-Ktext Kan^r vector digest included 1 µl pGEX-Ktext Kan^r vector, 2 µl 10X buffer, 0.5 µl PvuI, 0.5 µl ScaI, 1 µl H₂O. Both digests were incubated at 37 °C for 3 hours. Both the digested PCR fragment and the digested vector DNA were purified from 0.8% TAE agarose gels (Figure 1A).

- The PCR product was then ligated into Cst-04 vector. The ligation reaction contained 7 µl gel-purified Kan^r gene (cut with SacI/XbaI), 1 µl gel-purified Cst-04 vector (cut with BamHI/EcoRI), 1 µl 10X Ligation Buffer 1 µl T4 DNA ligase, and was incubated on ice overnight. The ligated PCR product was then transformed into the TG1 competent cells. The transformation reaction conditions included 500 µl (thawed on ice) TG1 Competent cells and 5µl pGEX-KT-exT-kan^r-CNS-CstI ligation rxn. The cell/DNA mixture was incubated on ice for 30 minutes, and the cells were heat shocked at 42°C for 45 seconds and then incubated on ice again for 2 minutes. 500 µl LB broth was added and the mixture was shaken at 37°C for 1 hour. The transformation reactions were then plated on LB Kan^r plates and

incubated 37°C overnight. The results of the transformation reactions are shown in Figure 1B.

Positive clones were screened for using the following method. Two milliliters of 9x LB/Kanamycin (10µg/ml) culture was incubated with individual transformants at 37°C o/n using 250 RPM shaking. 1.5 milliliters of the overnight culture was transferred to an eppendorf tube to isolate plasmid DNA using Wizard Mini-Prep Kit (Qiagen, Valencia, CA). An insert-containing colony (Cst-04-Kan5, Figure 1C) was expanded in 100 ml of LB culture in order to isolate more plasmid DNA.

The Cst-04-Kan5 plasmid-containing colony was screened for kanamycin and ampicillin resistance. Cst-04-Kan5 colony was streaked on both AFLB Kan^r and Amp^r plates, which were incubated overnight at 37 °C. Figure 1D shows that, in colony Cst-04Kan5, the kanamycin resistance gene is active and the Ampicillin resistance gene is inactive.

The Cst-04Kan5 plasmid was transformed into E. coli JM109 cells using 20 µl JM109 competent cells in 0.34 µl 1.42 M beta mercaptoethanol, incubated on ice for 10 minutes, then adding 1µl (100 ng) Cst-04-Kan5 plasmid. The cell/DNA mixture was incubated ice for 30 minutes, then heat shocked at 42 °C for 45 seconds. The reaction was then incubated on ice for 2 minutes, at which time 80 µl SOC or LB was added. The reaction mixture was shaken at 37°C for 1 hour, then plated on LB Kan^r plates. Identification and confirmation of the Cst-04-Kan5 plasmid DNA was carried out with a restriction enzyme digestion of plasmid DNA isolated from positive transformants, using NdeI, SalI, PstI restriction enzymes. The restriction fragment sizes were ~7.3 kb for one cut, such as NdeI or SalI. Three bands (~1.7kb, ~2.2kb, and ~3.2kb) were observed when Cst-04-Kan5 DNA was cut with PstI (See Figure 1C, lane 4).

Starter cultures of Cst-04Kan5 plasmid-containing cells were produced and used to inoculate 100ml cultures for the generation of cell lysates. Centrifuged cell pellets resulting from the large-scale cultures were resuspended in 5ml H₂O prior to lysis in a French press. The resultant lysate was centrifuged at 4°C, 18,000 RPM for 20 minutes. The clarified lysate was subsequently used for activity analysis and the remainder of the lysate was stored at -20°C.

The activity of the cell lysates was determined under the assay conditions illustrated in Table 1.

Table 1.

Reagent	Final Concentration	Stock Concentration	Amounts
CTP	1 mM	100 mM	1 μ l
NAN	1 mM	200 mM	0.5 μ l
LacPsy	0.5 mM	2.5 mM	20 μ l
MgCl ₂	50 mM	1M	5 μ l
Tris pH 8	100mM	1M	10 μ l
Lysate	15 %	Crude	15 μ l
dH ₂ O			63 μ l
Total reaction volume			100 μ l

Table 1 lists the reagents, and concentrations and volumes thereof, used in the lysate activity assays.

The lysate assay reagents were mixed and incubated at 37°C. Time points were taken at 0 minutes and 1 hour. A negative control (pGEX-KT-exT-kan^r vector without insert) was also included. All time points were analyzed using thin layer chromatography (CHCl₃:CH₃OH:H₂O:NH₄OH:60:40:5:1 respectively). The plates were air dried, dipped in anisaldehyde and heated on a hot plate until the spots developed. Cst-04Kan5 lysate was also assayed for activity using lacto-N-neotetraose as substrate. The lacto-N-neotetraose substrate activity is illustrated in Figure 1E. Activity of lysates from Cst-04Kan5 plasmid-containing cells was 8500 units/liter.

Example 2: Modification of the polylinker of pCWori+ Kan^r expression vector.

pCWori+ Kan^r (Cst-04Kan5) contains the genes encoding *N. meningitidis* CMP-NAN synthetase (CNS) and *Campylobacter jejuni* α 2,3 Sialyl Transferase (CstII) at the multiple cloning site. This vector appears to give high levels of expression of recombinant proteins but is hard to use due to the limited number of restriction sites in the multiple cloning site (MCS). Therefore, the multiple cloning site was modified and expanded as described herein.

The multiple cloning site starting at NdeI restriction site and extending to the start of the inactive Amp^r gene from Cst04Kan5 was generated using PCR to

generate cDNA with modified multiple cloning sites at 5' (Pewmcs (NdeI) - ATCGATCGACATATGGGATCCGAGCTCAAGCTTCTAGACTCGAGGAATT CCGGTACCGTCGACATCGATGATAAGCTGTCAAA) (SEQ ID NO:6) and 3' (ScaI-ATTCCAAGTAGTACTACTCTTCTTTTCAA) (SEQ ID NO:7) ends of pCWIN1 construct. The PCR primers set for pre pCWIN2 construct are 5' (Bgl II-CAATTATATAGATCTATCGATGCTTAGGAGGT) (SEQ ID NO:8) and 3' (CstI-Xba-TTGCCTTATTCTAGATCATTAGTGGTGATGGTGGTG) (SEQ ID NO:9). The PCR products were then subcloned into Cst04kan5 (pCWori+ kan^r-CNS-CstI) vector, transformed into TG1 cells, and screened for the correct construct.

Two PCR reactions were conducted, using 10 ng Cst04Kan5 cDNA as a template. The first reaction contained 1 µg (1 µl) Pewmcs/ScaI-pcw primer, 78 µl H₂O, 8 µl dNTP mixture, 10 µl 10X buffer, and 1 µl Vent polymerase. The second reaction contained 1 µg (1 µl) 5' pcBglIII/CstI-Xba primer, 78 µl H₂O, 8 µl dNTP mixture, 10 µl 10X buffer, and 1 µl Vent polymerase. The PCR reaction parameters included a 5 minute cycle at 95 °C, followed by the addition of 1 µl of Vent polymerase and thirty cycles of the following temperature pattern: 94 °C for one minute, 55 °C for one minute, 72 °C for one minute.

The PCR products were subjected to a restriction digest. The first PCR reaction product ("pCWIN1" insert) digest included 16 µl PCR rxn, 2 µl 10X buffer, 0.5 µl NdeI, 0.5 µl ScaI, 1 µl H₂O, and the second PCR reaction product ("pre pCWIN2" insert) digest included 16 µl PCR rxn, 2 µl 10X buffer, 0.5 µl BglIII, 0.5 µl EcoRI, 1 µl H₂O. A pCWori Kan^r Cst04Kan5 vector was prepared for insertion of the first PCR product by incubation of 2 µl (1 µg) Cst04Kan5 vector, 2 µl 10X buffer, 0.5 µl NdeI, 0.5 µl ScaI, 1 µl H₂O. A pCWori Kan^r Cst04Kan5 vector was similarly prepared for insertion of the second PCR product by incubation of 2 µl (1 µg) Cst04Kan5 vector, 2 µl 10X buffer, 0.5 µl BamHI, 0.5 µl EcoRI, 1 µl H₂O.

The digested PCR fragments and digested vectors were purified from 0.8% TAE agarose gels (Figure 2A). The PCR products were then subcloned into Cst-04kan5 vectors by ligation (Table 2) and electroporated into the TG1 competent cells. Electroporation reactions included 30 µl thawed (on ice) TG1/DH5α electrocompetent cells and 3 µl ligation reaction mixture. The DNA/cell electroporation mixture was transferred to a chilled cuvette, and the cells were subjected to electroporation using pulses of 2.5 KV, R5 resistance, and 129 ohms. 0.9

ml SOC media was then added to the reaction mixture, and the entire culture was incubated at 37 °C for one hour, at which time the electoporation product was incubated overnight after plating on LB agar plates containing 50 µg/ml Kan^r.

5 Table 2

10	1. pCWIN1	
	Gel-purified pCWIN1 insert (digested A)	7 µl
	Gel-purified Cst-04kan5 vector (digested B)	1 µl
	10X Ligation Buffer	1 µl
15	T4 DNA Ligase	1 µl
	2. pre pCWIN2	
	Gel-purified pre pCWIN2 insert (digested C)	7 µl
	Gel-purified Cst-04kan5 vector (digested D)	1 µl
20	10X Ligation Buffer	1 µl
	T4 DNA Ligase	1 µl
	3. pCWIN2	
	Gel-purified pCWIN1#5 insert (digested E)	7 µl
25	Gel-purified pre pCWIN2#11 vector (digested F)	1 µl
	10X Ligation Buffer	1 µl
	T4 DNA Ligase	1 µl

Table 2 illustrates the ligation reaction conditions for pCWIN1, pre-pCWIN2, and pCWIN2 PCR reaction products. Both pCWIN1 and pre pCWIN2 ligations were incubated at 4°C overnight.

Screening of transformants for positive clones was then conducted.

Five colonies pCWIN1-containing TG1 cells were selected, as were 9 colonies of pre-pCWin2 in DH5α, and 18 colonies of pCWin2 in TG1. Each was placed into 2 ml TB/kanamycin (50µg/ml) and incubated at 37°C for 5 hours with shaking at 250 RPM. 1.5 milliliters of each culture was transferred to an eppendorf tube to isolate plasmid DNA using Wizard Plus Mini-Prep Kit (Qiagen, Valencia, CA). Each plasmid DNA preparation was subjected to restriction digestion with the appropriate restriction endonucleases as described above. The digestion reactions were then analyzed on agarose/TAE gels (Figures 2B – 2D).

One colony of pCWin1 (colony #5) and one colony of pre pCWin2 (colony #11) contained the appropriate sized inserts. Specifically, the expected size for the pCWin1 insert was 5 kb and the expected size for the pre-pCWin2 insert was 7 kb (Figure 2B). pCWin1 and pre pCWin2 plasmid DNA were digested with NdeI and

- ScaI. NdeI and ScaI digestion of pCWin1 plasmid generated bands of 750 bp and 4.2kb, and the same digestion of pre-pCWin2 plasmid generated bands of 2.8 kb and 4.2 kb (Figure 2C). Figure 2D illustrates the result of ligating the 750 bp of pCWin1 with the 4.2 kb fragment of pre-pCWin2 to generate the pCWin2 expression vector.
- 5 The difference between pCWin1 and pCWin2 expression is that pCWin1 has two BamHI sites, one in being in the tac promoter and the other being in the multiple cloning site (down stream of NdeI). pCWin2 has only one BamHI site, in the multiple cloning site, and the BamHI that resided in the tac promoter was destroyed.
- 10 Example 3: Addition of Maltose Binding Protein to pCWin2 Kan^r Expression Vector

The *E.coli* *malE* gene, encoding a maltose binding protein, was subcloned into the pCWin2 kan^r bacterial expression vector. The *malE* gene was PCR amplified from pMal-c2X, ligated into the multiple cloning site of pCWin2 kan^r, and subsequently transformed into electrocompetent DH5a *E. coli*. The final product, a pCWin2MBP kan^r bacterial fusion tag expression vector, was created as described below.

Restriction endonuclease digestion of pCWin2 kan^r and pMAL-c2X amp^r was conducted to prepare the *malE* maltose binding protein cDNA and the pCWin2 vector cDNA for insertion of the *malE* cDNA into the pCWin2 vector. Digestion of the *malE* cDNA was conducted using 2 µl of pMAL -c2X vector DNA (1µg/µl), 2 µl 10X BamHI NEbuffer, 2 µl 10X purified BSA, 1 µl NdeI, 1 µl BamHI, and 12 µl dH₂O. Digestion of the vector was conducted using 2 µl pCWin2 vector DNA 0.8µg/µl, 2 µl 10X BamHI NEbuffer, 2 µl 10X purified BSA, 1 µl NdeI, 1 µl BamHI, and 12 µl dH₂O.

The restriction enzyme digestions were incubated at 37°C for two hours. The reactions were stopped by adding 3 µl Blue/Orange 6x Agarose Loading Dye. The digestions were then loaded onto separate 0.7% agarose/TAE gels, and electrophoresed at 135 volts until the dyes migrated to the lower third of the gel. An image of the pCWin2 vector digestion agarose gel was captured using a digital camera (Figure 3B). An image of the polyacrylamide gel containing the purified product from the digestion of *malE* is shown in Figure 3A.

The linearized pCWin2 kan^r and *malE* fragments were gel purified. Using the UV box to illuminate the DNA, the bands of DNA were excised using a

sterile scalpel. The pCWin2 kan^r DNA was approximately 5 kb, and the *malE* gene was approximately 1.2 kb. The excised agarose wedges were placed into Ultrafree DA agarose extraction spinfilters (Millipore, Bellerica, MA), and microcentrifuged at 10,000 xg for 5 minutes. The filtrates were transferred to YM-100 spinfilters (Millipore, Bellerica, MA), and the DNA was washed by adding 300 μ l dH₂O. The spinfilters were centrifuged at 500 xg for 15 minutes, and the wash step was repeated two additional times. The last wash step concentrated the DNA to an approximate volume of 25 μ l, at which time the column was inverted into another autoclaved 1.7 mL eppendorf tube. The DNA retentate was collected by microcentrifuging the eppendorf at 1000xg for one minute.

The ligation of gel purified *malE* and linearized pCWin2 kan^r was performed in an autoclaved 0.5 mL eppendorf microcentrifuge tube. The ligation reaction included 7 μ l of purified *malE* DNA that was digested with NdeI and BamHI, 1 μ l of linearized pCWin2 kan^r that was digested with NdeI and BamHI, 1 μ l 10X ligase buffer, 1 μ l T4 DNA ligase. The ligation reaction was incubated at room temperature for three hours. In the vector control ligation reaction, dH₂O was substituted for the *malE* DNA.

The ligation reactions were transformed into electrocompetent DH5 α *E.coli*. After a three hour ligation incubation, one aliquot of electrocompetent DH5 α *E.coli* was removed from a -81°C freezer, and placed on ice to thaw. 20 μ l of the cells was aliquoted into chilled, autoclaved 1.7 mL microcentrifuge tubes, and then one microliter from each ligation reaction was added to the cells. Immediately, the reactions were transferred to chilled electroporation cuvettes. The cells were electroporated with a 2.5kV 6 msec pulse as described in the manufacturer's instructions. Then one milliliter of AFLB SOC media was added to the transformation reactions, and the entire volume was transferred to an autoclaved 1.7 mL microcentrifuge tube. The transformation reactions were incubated at 37°C for one hour with shaking at 250 rpm. After incubating the cells for an hour, 100 μ l from each transformation reaction was plated by spreading onto LB agar kan^r plates. The plates were incubated at 37°C overnight.

Results from the ligation and transformation reactions are as follows. The pCWin2 MBP vector plating resulted in thirteen colonies, and out of ten colonies selected, nine were positive for the recombinant vector. The pCWin2 vector control plating did not contain any colonies. Colonies from the pCWin2 MBP vector LB agar

plates were selected and used to inoculate 2 ml AFLB kan^r. The starter cultures were grown overnight at 37°C, with shaking at 250 rpm. Plasmid DNA was isolated from the transformants and screened for the correct *malE* insert by a double digestion with NdeI and BamHI restriction endonucleases.

- 5 Restriction digestion of miniprep DNA was conducted in a reaction mixture containing 12 µl Miniprep DNA, 1 µl NdeI endonuclease, 1 µl BamHI endonuclease, 1.5 µl 10XBamHI NEBuffer, and 1 µl 10X purified BSA. The digestion reactions were mixed and incubated at 37°C for one hour. After incubation, 3 µl of 6x Agarose Gel Loading Dye was added to each restriction digestion. The
10 restriction digestions were then loaded into the wells of a 0.7% agarose/TAE gel. The samples were electrophoresed at 135 volts until the dye migrated to the lower third of the gel. The gel was then removed from the gel box, and the image captured with a digital camera (Figure 3C).

- Large scale purified pCWin2MBP vector DNA was isolated from
15 transformant #1 using the HiSpeed Plasmid Maxi Kit (Qiagen, Valencia, CA). A 2 mL AFLB kan^r starter culture was inoculated with 10 µl pCWin2MBP DH5a *E.coli* overnight culture. This starter culture was grown overnight in a 37°C incubator, with shaking at 250 rpm. The overnight starter culture was used to inoculate two 125 mL
AFLB kan^r cultures, and these larger scale preps were grown overnight at 37°C with
20 shaking at 250 rpm. DNA from the large scale preparation was used for sequencing of the *malE* insert subcloned between the NdeI and BamHI restriction sites (MWG Biotech, High Point, NC). The sequence of the vector is set forth in SEQ ID NO:3.

Example 4: Preparation and Characterization of pCWin2-MBP-SBD₃₉(pMS₃₉) Vector

- 25 The pMS₃₉ Kan^R expression vector was created from the pCWIN2-MBP- SBD-ST3 Gal III (GalB1,3(4)GlcNAc a2,3-Sialyltransferase) Δ73 construct, removing the the ST3 Gal III Δ73 gene and replacing it with the Multiple Cloning site (MCS) from the pcWIN2 vector. Selection was of final construct was determined by restriction enzyme analysis with Nde I and Xba I (there is no Xba I site the pCWIN2-
30 MBP-SBD ST3 Gal III construct) digestion and sequence confirmation. The final construct was designated the pCWin2-MBP-SBD (pMS₃₉) Kan^r expression vector. The several steps of the preparation of this vector are detailed in Figures 7A-7G.

Example 5: Preparation and Characterization of the pCWin2-MBP-MCS-SBD (pMXS₃₉) Vector

The pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector was constructed according to the following method. The Starch Binding Domain (SBD) insert was
5 isolated by PCR using the 5' primer (XhoI-SBD-39-5'
TGTATCCTCGAGATTGTGGCGACCGCGGCACCAC) (SEQ ID NO:12) and the
3' primer (3' SalI-AAGCTTGTCTGACTCATTAGCGCCAGGTATCGGTCACGG)
(SEQ ID NO:13). The PCR products were gel purified and ligated into PCR-Blunt
vector. The correct SBD insert (in the PCR-Blunt vector) was digested with XhoI and
10 SalI, subcloned into XhoI-SalI digested pCWin2-MBP kan^r vector, transformed into
TB1 cells and screened for the correct construct. The several steps of the preparation
of this vector are detailed in Figures 8A-8E.

The disclosures of each and every patent, patent application, and publication
15 cited herein are hereby incorporated herein by reference in their entirety. While the
invention has been disclosed with reference to specific embodiments, it is apparent
that other embodiments and variations of this invention may be devised by others
skilled in the art without departing from the true spirit and scope of the invention.
The appended claims are intended to be construed to include all such embodiments
20 and equivalent variations.

CLAIMS

What is Claimed is:

- 5 1. A method of providing a therapeutic protein to a customer, said method comprising cloning a nucleic acid encoding said protein into a pCWin1 expression vector as set forth in SEQ ID NO:1, expressing said protein therefrom, and providing said protein to said customer.
- 10 2. A method of providing a therapeutic protein to a customer, said method comprising cloning a nucleic acid encoding said protein into a pCWin2 expression vector as set forth in SEQ ID NO:2, expressing said protein therefrom, and providing said protein to said customer.
- 15 3. A method of providing a therapeutic protein to a customer, said method comprising cloning a nucleic acid encoding said protein into a nucleic acid vector selected from the group consisting of:
- a) a pCWin2/MBP expression vector as set forth in SEQ ID NO:3;
- b) a pCWin2-MBP-SBD (pMS₃₉) expression vector as set forth in
- 20 SEQ ID NO:10; and
- c) a pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector as set forth in SEQ ID NO:11;
- expressing said protein therefrom, and providing said protein to said customer.
- 25 4. The method of claim 3, wherein said nucleic acid vector comprises a protease cleavage site coding sequence at a location selected from the group consisting of:
- a) between the MBP coding sequence and the therapeutic protein
- 30 coding sequence; and
- b) immediately prior to the start of the C-terminus of the MBP coding sequence.

5. The method of claim 2 or 3, wherein said protein is selected from the group consisting of erythropoietin, human growth hormone, granulocyte colony stimulating factor, interferons alpha, -beta, and -gamma, Factor IX, follicle stimulating hormone, interleukin-2, erythropoietin, anti-TNF-alpha, and a lysosomal hydrolase.

6. The method of claim 5, wherein said lysosomal hydrolase is selected from the group consisting of beta-glucosidase, alpha-galactosidase-A, beta-hexosaminidase, beta-galactosidase, alpha-galactosidase, alpha-mannosidase, beta-mannosidase, alpha-L-fucosidase, beta-glucuronidase, alpha-glucosidase, alpha-N-acetylgalactosaminidase, and acid phosphatase.

7. A method of providing a protein to a customer, said method comprising cloning a nucleic acid encoding said protein into a pCWin1 expression vector as set forth in SEQ ID NO:1, expressing said protein therefrom, and providing said protein to said customer.

8. A method of providing a protein to a customer, said method comprising cloning a nucleic acid encoding said protein into a pCWin2 expression vector as set forth in SEQ ID NO:2, expressing said protein therefrom, and providing said protein to said customer.

9. A method of providing a protein to a customer, said method comprising cloning a nucleic acid encoding said protein into a nucleic acid vector selected from the group consisting of:

- a) a pCWin2-MBP expression vector as set forth in SEQ ID NO:3;
 - b) a pCWin2-MBP-SBD (pMS₃₉) expression vector as set forth in SEQ ID NO:10; and
 - c) a pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector as set forth in SEQ ID NO:11;
- expressing said protein therefrom, and providing said protein to said customer.

10. The method of claim 7, 8 or 9, wherein said protein is selected from the group consisting of a glycosyltransferase and a sugar nucleotide-generating enzyme.

5 11. A method of providing a protein to a customer, said method comprising providing a pCWin1 vector as set forth in SEQ ID NO:1 to a protein production facility, wherein a nucleic acid encoding said protein is cloned into said expression vector and said protein is expressed therefrom in said protein production facility, and providing said protein to said customer.

10

12. A method of providing a protein to a customer, said method comprising providing a pCWin2 vector as set forth in SEQ ID NO:2 to a protein production facility, wherein a nucleic acid encoding said protein is cloned into said expression vector and said protein is expressed therefrom in said protein production facility, and providing said protein to said customer.

15

13. A method of providing a protein to a customer, said method comprising providing a nucleic acid vector selected from the group consisting of:

- 20 a) a pCWin2-MBP expression vector as set forth in SEQ ID NO:3;
b) a pCWin2-MBP-SBD (pMS₃₉) expression vector as set forth in SEQ ID NO:10; and
c) a pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector as set forth in SEQ ID NO:11;

25 to a protein production facility, wherein a nucleic acid encoding said protein is cloned into said expression vector and said protein is expressed therefrom in said protein production facility, and providing said protein to said customer.

14. The method of claim 2, 3, 4, 7, 8 or 9, wherein said method further comprises prior to providing said protein to said customer, at least one glycosyl moiety is added to said protein.

30

15. The method of claim 14, wherein said glycosyl moiety is added to said protein in vitro.

16. A method of providing a protein to a customer, said method comprising cloning a nucleic acid encoding said protein into nucleic acid vector selected from the group consisting of:

- a) a pCWin1 vector as set forth in SEQ ID NO:1;
- b) a pCWin2 vector as set forth in SEQ ID NO:2;
- c) a pCWin2-MBP vector as set forth in SEQ ID NO:3;
- d) a pCWin2-MBP-SBD (pMS₃₉) vector as set forth in SEQ ID NO:10; and
- e) a pCWin2-MBP-MCS-SBD (pMXS₃₉) vector as set forth in SEQ ID NO:11;

further wherein said method comprises inserting said vector into a bacterial host cell, expressing said protein in said host cell, and providing said protein to said customer.

17. The method of claim 16, wherein said method further comprises prior to providing said protein to said customer, at least one glycosyl moiety is added to said protein.

18. The method of claim 16, wherein said glycosyl moiety is added to said protein in vitro.

19. The method of claim 16, wherein said expression vector further comprises an affinity tag coding sequence.

20. An isolated pcWIN1 expression vector comprising the sequence set forth in SEQ ID NO:1.

21. An isolated pcWIN1 expression vector consisting of the sequence set forth in SEQ ID NO:1.

22. An isolated pcWIN2 expression vector comprising the sequence set forth in SEQ ID NO:2.

23. An isolated pcWIN2 expression vector consisting of the sequence set forth in SEQ ID NO:2.

24. An isolated pcWIN2/MBP expression vector comprising the sequence set forth in SEQ ID NO:3.

25. An isolated pcWIN2/MBP expression vector consisting of the sequence set forth in SEQ ID NO:3.

26. The pcWIN2/MBP expression vector of claim 24, wherein the pCWIN2/MBP vector comprises a protease cleavage site coding sequence adjacent to the MBP coding sequence.

27. An isolated pCWin2-MBP-SBD (pMS₃₉) vector comprising the sequence set forth in SEQ ID NO:10.

28. An isolated pCWin2-MBP-SBD (pMS₃₉) vector consisting of the sequence set forth in SEQ ID NO:10.

29. An isolated pCWin2-MBP-MCS-SBD (pMXS₃₉) vector comprising the sequence set forth in SEQ ID NO:11.

30. An isolated pCWin2-MBP-MCS-SBD (pMXS₃₉) vector consisting of the sequence set forth in SEQ ID NO:11.

31. The pCWin2-MBP-SBD (pMS₃₉) expression vector of claim 27, wherein the pCWin2-MBP-SBD (pMS₃₉) vector comprises a protease cleavage site coding sequence immediately prior to the start of the C-terminus of the MBP coding sequence.

32. A method of expressing a protein, said method comprising cloning a nucleic acid encoding said protein into a pCWin1 expression vector as set forth in SEQ ID NO:1 and expressing said protein therefrom.

33. A method of expressing a protein, said method comprising cloning a nucleic acid encoding said protein into a pCWin2 expression vector as set forth in SEQ ID NO:2 and expressing said protein therefrom.

- 5 34. A method of expressing a protein, said method comprising cloning a nucleic acid encoding said protein into a nucleic acid vector selected from the group consisting of:
- a) a pCWin2/MBP expression vector as set forth in SEQ ID NO:3;
 - b) a pCWin2-MBP-SBD (pMS₃₉) expression vector as set forth in
 - 10 SEQ ID NO:10; and
 - c) a pCWin2-MBP-MCS-SBD (pMXS₃₉) expression vector as set forth in SEQ ID NO:11;
- and expressing said protein therefrom.
- 15 35. The method of any one of claims 32-34, wherein said protein is expressed in a prokaryotic cell.

1/22




FIG. 1A




FIG. 1B

2/22



FIG. 1C




FIG. 1D

3/22




FIG. 1E

4/22




FIG. 2A




FIG. 2B

5/22




FIG. 2C




FIG. 2D

6/22

λ std
malE




FIG. 3A

1 kb pCWin2
std NdeI/BamHI




FIG. 3B




FIG. 3C

8/22

gcacgtggtgtcacgctcgctgttggatggcttcattcagctccggttcccaacgatca
aggcgagttacatgatcccccatgttgtgcaaaaaagcggttagctccttcggctcctccgat
cgggggggggggaaagccacgttgtgtctcaaaatctctgatgttacattgcacaagataa
aaatatatcatcatgaacaataaaactgtctgtctacataaacagtaataacaaggggtgta
tgagccatattcaacgggaaacgtcttgcctcaggccgcgattaaattccaacatggatgct
gatttatattgggtataaaatgggctcgcgataatgtcgggcaatcagggtgcgacaatctatcg
actgtatgggaagcccgatgcgcagagttgtttctgaaacatggcaaaaggtagcgttgcca
atgatgttacagatgagatggctcagactaaactggctgacggaatttatgcctcttccgacc
atcaagcattttatccgtaactcctgatgatgcatgggtactcaccactgcgatccccgggaa
aacagcattccagggtattagaagaatatcctgattcagggtgaaaatattgttgatgcgctgg
cagtggtcctgcgcgggttgcatctcattcctgtttgttaattgtccttttaacagcgatcgc
gtatttcgtctcgcctcaggcgcaatcacgaatgaataacgggttggttgatgcgaagtgaatt
tgatgacgagcgtaattggctggcctgttgaaacagctcggaaagaaatgcataagctattgc
cattctcacggattcagtcgctcactcatgggtgatttctcacttgataaccttatttttgac
gaggggaaattaataggttgatttgatgttgacgagtcggaatcgcagacogataaccagga
tcttgccatcctatggaaactgcctcgggtgagttttctccttcattacagaaacggctttttc
aaaaatatggtattgataatcctgatatgaataaaatgcagtttcatattgatgcctgatgag
tttttctaaagtaactactcctttttcaatattattgaagcatttatcagggttattgtc
tcatgagcggatacatatttgaaatgtatttagaaaaataaacaataagggggtccgcgcaca
tttccccgaaaagtgcacctgaogatgaaattgttaaacgttaatatatttgtaaatttcgc
gttaaatttttgttaaatcagctcattttttaaccaataggccgaaatcggcaaaatccctt
ataaatcaaaagaatagcccgagatagggttgagtggtgttccagtttggaacaagagtgca
ctattaaagaacgtggactccaacgtcaaaggcgcaaaaacogtctatcaggggcgatggccc
actacgtgaaccatcacccaatcaagtttttggggctcaggtgcggtaaagctctaaatc
ggaaocctaaaggagcccccgatttagagcttgacgggggaaagccggcgaaacgtggcgaga
aaggaaagggaagaaagcgaaaggagcgggcgctagggcgctggcaagtgtagcggctcacgct
gcgcgtaaccaccaccccgccgcgcttaatgcgcgcgtacaggggcgctactatgggtgct
ttgaogcatcgtctaagaaaccattattatcatgacattaacctataaaaaataggcgatca
cgaggccctttcgtcttcaagcagatctgaaaaaaaagcccgctcattaggcgggctcagat
ctgctcatgtttgacagcttatcatcgatgtcgacggtaccgaattcctcgagctctagaaag
cttgagctcggatcccatatgacctcctaagcatcgatggatcctgtttcctgtgtgaaatt

Fig. 4A

9/22

gttatccgctcacaattccacacattatatacgagccgatgattaattgtcaacagggggatgg
ggagtaagctgatcctgtttcctgtgtgaaattgttatccgctcacaattccacacattata
cgagccgatgattaattgtcaacagggggatggggagtaagctcatcgatggatcgatcctg
tttctgtgtgaaattgttatccgctcacaattccacacattatatacgagccggaagcataaa
gtgtaagcctgggtgacctaatgagtgagctaaacttacattaattgcggtgcgctcactgc
ccgctttccagtcgggaaacctgtcgtgccaggacaccatcgaaatgggtgcaaaacctttcgc
ggataggcatgatagcgcccggaagagagtgcaattcagggtggatgaaaccagtaac
gttatacgatgtcgcagagtatgccggtgtctcttatcagaccgtttcccgctggatgaacc
aggccagccacgtttctgcgaaaacgcgggaaaaagtgaagcggcgatggcggagctgaat
tacattcccaaccgctgggcacaacaactggcgggcaaacagtcgttgcgtgattggcgttgc
cacctccagtcctggccctgcacgcgcgtgcgcaaatgtgcggcgatataatctcgcgcgc
atcaactgggtgccagcgtgggtgtgcgatggtagaacgaagcggcgatgaagcctgtaaa
gcggcggtgcacaatcttctcgcgcaacgcggtcagtgggctgatacctaactatccgctgga
tgaccaggatgccattgtcgtggaagctgctgcactaatgttccggcgttatttcttgatg
tctctgaccagacacccatcaacagtgatttttctcccatgaagacggatcgcgactgggc
gtggagcatctggctgcattgggtcaccagcaaatcgcgctgttagcgggcccattaaagtct
tgtctcggcgcgtctgcgtctggctggctggcataaatactcactcgcaatcaaatcagc
cgatagcgggaacgggaagcgcagctggagtgccatgtccggttttcaacaaacctgcgaatg
ctgaatgagggcatcgttcccactgcgatgctggttgccaaagatcagatggcgctgggcgc
aatgocgcgcattaccagagtcgggctgcgcgttggtgcggatatctcggtagtgggatacg
acgataccgaagacagctcatgttatacccgccgttaaccaccatcaaacaggattttcgc
ctgctggggcaaacagcgtggaccgcttgctgcaactctctcagggccaggcggatgaagg
caatcagctgttgccgctctcactggtgaaaagaaaaaccacctggcgcccaatacgcaaa
ccgctctcccgcgcggttgccgattcattaatgcagctggcacgacaggtttccgactg
gaaagcggcagtgagcgcaacgaattaatgtaagttagctcactcattaggcaccaccagg
ctttacactttatgcttcgggctcgatggcggtttcgggtgatgacgggtgaaaacctctgaca
catgcagctcccgagacggtcacagcttgtctgtaagcggatgcggggagcagacaagccc
gtcagggcgcgctcagcgggtgttgccgggtgtcggggcgacgcatgaccagtcacgtagc
gatagcggagtgatactggcttaactatgcggcatcagagcagattgtactgagagtgacac
cattatgocggtgtgaaataccgcacagatgcgtaaggagaaaaataccgcacagcgctctt
ccgcttctcgcgtcactgactcgcgtcgcgtcggctcgttcgggtgcggcgagcgggtatcagct

FIG. 4B

10/22

cactcaaaggcggttaatacggttatccacagaatcaggggataacgcaggaaagaacatgtg
agcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgcttgctggcgtttttccata
ggctccgccccctgacgagcatcacaaaaatcgacgctcaagtcagagggtggcgaaacccg
acaggactataaagataaccaggcgtttccccctggaagctccctcgtgcgctctcctgttcc
gacctgcgcgttacccgataacctgtccgcctttctcccttcgggaagcgtggcgctttctc
atagctcacgctgtaggatctcagttcgggtgtaggtcgttcgctccaagctgggctgtgtg
cacgaaccccccggttcagccgaccgctgcgccttatccggtaactatcgtcttgagtccaa
cccggttaagacacgacttatcgccactggcagcagccactggtaacaggattagcagagcga
ggatatgtaggcgggtgctacagagttcttgaagtgggtggcctaactacggctacactagaagg
acagtatttggtatctgcgctctgctgaagccagttaccttcggaaaaagagttggtagctc
ttgatccggcaaaacaaaccgcgtggtagcgggggttttttggtttgcaagcagcagatta
cgcgcaaaaaaaaggatctcaagaagatcctttgatcttttctacggggtctgacgctcag
tggaacgaaaaactcacgttaagggattttgggtcatgagattatcaaaaaggatcttcacct
gatccttttaaatataaaatgaagttttaaatcaatctaaagtatatatgagtaaacttgg
ctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcgttca
tccatagttgcctgactccccgtcgtgtagataactacgatacgggagggcttaccatctgg
cccagtgctgcaatgatacogcgagaccacgctcaccggctccagatttatcagcaataa
accagccagccggaaggccgagcgcagaagtggctcctgcaactttatccgcctccatccag
tctattaattggtgcgggaagctagagtaagtagttcgccagttaatagtttgcgcaacgt
tggtgccattgctgcag

FIG. 4C

11/22

gcacgcgtgggtgtcacgcctcgtcgttttggtatggettcattcagctccgggtcccaacgatca
aggcgagttacatgatcccccatgtttgtgcaaaaaagcggttagctcctcgtgtcctccgat
cgggggggggggaaagccacgtttgtgtctcaaaatctctgatgttacattgcacaagataa
aaatatatcatcatgaacaataaaaactgtctgcttacataaaacagtaatacaagggtgtta
tgagccatattcaacgggaaacgtcttgctccaggccgcgattaaattccaacatggatgct
gatttatatgggtataaatgggctcgcgataatgtcgggcaatcaggtgcgacaatctatcg
actgtatgggaagcccgatgcgccagagttgtttctgaaacatggcaaggtagcgttgcca
atgatgttacagatgagatggctcagactaaaactggctgacggaatttatgcctcttccgacc
atcaagcattttatccgtactcctgatgatgcaggttactcaccactgcgatccccgggaa
aacagcattccaggtattagaagaatatcctgattcaggtgaaaaatattgttgatgcgctgy
cagtggtcctcgcgcgggttgattcgtattcctgtttgttaattgtccttttaacagcgatcgc
gtatttctcgtctcgtcaggcgcaatcacgaatgaataacgggtttgggtgatgcgagtgattt
tgatgacgagcgttaattggctggcctgttgaaacagctctggaagaaatgcataagctattgc
cattctcacccgattcagtcgtcactcatgggtgatttctcacttgataaccttatttttgac
gaggggaaattaatagggttgattgatgttgacgagtcggaatcgcgacccgataaccagga
tcttgccatcctatggaactgcctcggtagttttctccttcattacagaaacggctttttc
aaaaatatggtattgataaatcctgatatgaataaaattgcagtttcatttgatgctcgatgag
tttttctaaagtactactcttctttttcaatatatttgaaagcatttatcagggttattgtc
tcatgagcggatacatatttgaaatgtatttagaaaaataaaacaaatagggttccgcgcaca
tttccccgaaaagtgccacctgacgatgaaattgttaaacgttaatatattttgttaaaattcgc
gttaaatttttgttaaatcagctcattttttaaccaataggccgaaatcggcaaaatccctt
ataaatcaaaagaatagcccgagatagggttgagtgttggtccagtttggaacaagagtcca
ctattaaagaacgtggactccaacgtcaaaggcgcaaaaacgtctatcagggcgatggccc
actacgtgaaccatcacccaaatcaagttttttggggtcgagggtccgtaaaagctctaaatc
ggaaccctaaaggagcccccgatttagagcttgacggggaagccggcgaaacgtggcgaga
aagggaagggaagaaagcgaaaggagcgggcgctaggcgcgctggcaagtgtagcgtcacgct
gcgcgttaaccaccacacccgcgcgcttaatgcgcgcgtacaggcgcgctactatgggtgct
ttgacgcctcgtctaaagaaccattattatcatgacattaacctataaaaaataggcgatca
cgaggccctttcgtcttcaagcagatctgaaaaaaaaagcccgctcattaggcgggctcagat
ctgctcatgtttgacagcttatcatcgatgtcgacggtaccgaattcctcgagctagaaaag
cttgagctcggatcccatatgacctcctaagcatcgatagatcctgtttcctgtgtgaaatt

FIG. 5A

12/22

gttatccgctcacaattccacacattatacagagccgatgattaattgtcaacagggggatgg
ggagtaagctgatcctgtttcctgtgtgaaattgttatccgctcacaattccacacattata
cgagccgatgattaattgtcaacagggggatggggagtaagctcatcgatggatcgatcctg
tttctctgtgtgaaattgttatccgctcacaattccacacattatacagagccggaagcataaa
gtgtaaacgctgggggtgcctaattgagttagctaaacttacattaattgcgttgcgctcactgc
ccgctttccagtcgggaacctgtcgtgccaggacaccatcgaaatggtgcaaaccttttcgc
ggtatggcatgatagcgcccggaagagagtcaattcagggtggtgaatgtgaaaccagtaac
gttatcagatgtcgcagagtatgccggtgtctcttatcagaccgtttcccgctggtgaacc
aggccagccacgtttctgcgaaaaacgcgggaaaaagtgaagcggcgatggcgagctgaat
tacattcccaaccgcgtggcacaacaactggcgggcaaacagtcgttgctgattggcgttgc
cacctccagtcctggccctgcacgcgcgctgcgaaattgtcgcggcgattaaatctcgcgcg
atcaactgggtgccagcgtggtggtgtcgtatggtagaacgaagcgcgctcgaagcctgtaaa
gcggcggtgcacaatcttctgcgcgaacgcgctcagtgggctgatacattaactatccgcctgga
tgaccaggatgccattgctgtggaagctgcctgcactaatgttcggcggttatctcttgatg
tctctgaccagacacccatcaacagattattttctcccatgaagacggtagcgcactgggc
gtggagcatctggctgcattgggtcaccagcaaatcgcgctgttagcggggcccatagttc
tgtctcggcgctctgcgtctggctggctggcataaatatctcactcgcaatcaaatcagc
cgatagcggaaacgggaagcgcgactggagtgcattgtccggttttcaacaaaccatgcaaatg
ctgaatgagggcatcgttccactgcgatgctggttgccaacgatcagatggcgctgggcgc
aatgcgcgcattaccagctccgggctgcgcttggtgcggatatctcggtagtgggatacg
acgataccgaagacagctcatgttatatccgcgcttaaccacatcaaacaggatttttcgc
ctgctggggcacaaccagcgtggaccgcttgctgcaactctctcaggggcaggcggtgaaggg
caatcagctgttgccgctctcactgggtgaaaagaaaaaccacctggcgcccaatacgcaaa
ccgctctcccgcgcggttgccgattcattaatgcagctggcacgacaggtttcccgactg
gaaagcgggcagtgagcgcaacgcaattaatgtaagttagctcactcattaggcaccocagg
ctttacactttatgcttcgggctcgtatggcgcttcgggtgatgacggtgaaaacctctgaca
catgcagctcccgagacggtcacagcttgtctgtaagcggatgcggggagcagacaagccc
gtcagggcgcgctcagcgggtgttgccgggtgtcggggcgagccatgaccagtcacgtagc
gatagcggagtgtatactggcttaactatgcggcatcagagcagattgtactgagagtgcac
cattatgcggtgtgaaataccgcacagatgcgtaaggagaaaaataccgcacagcgctctt
ccgcttctcgcctcactgactcgcgtcgctcggttcggctgcggcgagcgggtatcagct

FIG. 5B

13/22

cactcaaaggcggttaatacggttatccacagaatcaggggataaacgcaggaaagaacatgtg
agcaaaaggccagcaaaaggccaggaaaccgtaaaaaggccgcgttgcgtggcgtttttccata
ggctccgccccccctgacgagcatcacaaaaatcgacgctcaagtcagaggtggcgaaaccg
acaggactataaagataaccaggcgtttccccctggaagctccctcgtgcgtctctcgtttcc
gacctgcgcgttacccggataacctgtccgcctttctcccttcgggaagcgtggcgctttctc
atagctcacgctgtaggtatctcagttcgggtgtaggtcgttcgctccaagctgggctgtgtg
cacgaaccccccggttcagcccgaccgctgcgccttatccggtaactatcgtcttgagtccaa
cccgtaagacacgacttatcgccactggcagcagccactggtaacaggattagcagagcga
ggtatgtaggcgggtgctacagagttcttgaagtgggtggcctaactacggctacactagaagg
acagtatttgggtatctgcgctctgctgaagccagttaccttcggaaaaagagttggtagctc
ttgatccggcaaaacaaaccaccgctggtagcgtggttttttggtttgcaagcagcagatta
cgcgagaaaaaaaggatctcaagaagatcctttgatctttttctacggggtctgacgctcag
tggaacgaaaaactcacgttaagggtattttggatcatgagattatcaaaaaggatcttcacct
gatccttttaaatataaaatgaagttttaaatcaatctaaagtatatatgagtaaaacttggt
ctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcgttca
tccatagttgcctgactccccgctgtagataactacgatacgggagggttaccatctgg
ccccagtgctgcaatgataaccgcgagaccacgctcacgggtccagatttatcagcaataa
accagccagccggaaggccgagcgcagaagtggctcctgcaactttatccgcctccatccag
totattaattgttgcgggaagctagagtaagtagttcgcaggttaatagtttgcgcaacgt
tgttgccattgctgcag

FIG. 5C

14/22

gcacgcgtggtgtcacgcctgcgtgttggtatggcttcattcagctccgggtcccaacgatca
 aggcgagttacatgatccccatgttgtgcaaaaaagcgggttagctccttcgggtcctccgat
 cgggggggggggaaagccacgctgtgtctcaaaatctctgatgttacattgcacaagataa
 aaatatatcatcatgaacaataaaactgtctgtctacataaacagtaataacaaggggtgtta
 tgagccatattcaacgggaaacgtcttgcctcaggccgcgattaaattccaacatggatgct
 gatttatatgggtataaaatgggctcgcgataatgtcgggcaatcagggtgcgacaatctatcg
 actgtatgggaagcccgatgcgcagagttgtttctgaaacatggc aaaggtagcgttgcca
 atgatgttacagatgagatggtcagactaaactggctgacggaatttatgcctcttccgacc
 atcaagcattttatccgtactcctgatgatgcattgttactcaccactgcgatccccgggaa
 aacagcattccaggtattagaagaatatcctgattcagggtgaaatatgttgatgcgtgg
 cagtggtcctgcgcgggttgcatctgatctcctgtttgtaattgtccttttaacagcgatcgc
 gtatttctgtctgcgtcaggcgcgaatcacgaatgaataacgggttggttgatgcgaagtattt
 tgatgacgagcgtaatggctggcctgttgaaacaagtctggaagaagtgcataagctattgc
 cattctcacgggatcagtcgtcactcatggtgatttctcacttgataacottatttttgac
 gaggggaaataataggttgatttgatgttggacgagtcggaatcgcagaccgataaccagga
 tcttgccatcctatggaactgcctcggtagttttctcctcattacagaaacggccttttctc
 aaaaatatggtatttgataatcctgatataaataatgacagtttctatttgatgctcgatgag
 tttttctaaagtaactactcttctcttttcaatatatttgaaagcattttatcagggttattgtc
 tcatgagcgggatacatatttgaaatgtatttgaaaaataaaacaaatagggttccgcgcaca
 tttcccccgaagtgccacctgacgatgaaattgtaaacgttaatatatttggttaaaattcgc
 gttaaatttttgttaaatcagctcattttttaaccaataggccgaaatcggcgaataatccctt
 ataaatcaaaagaatagcccgagatagggttgagtggtgttccagtttggaacaagagtcca
 ctattaaagaacgtggactccaacgtcaaaagggcgaaaaaccgtctatcagggcgatggccc
 actaogtgaaccatcaccccaaatcaagttttttggggctcgaggtgcgtaaaagctctaaatc
 ggaacctaaaggagcccccgatatttagagcttgacggggaaagccggcgaaacgtggcgaga
 aaggaagggaagaagcgaaaggagcgggcgctagggcgctggcaagtgtagcggctcacgct
 gcgcgtaaccaccacaccgcgcgcgttaatgcccgcgtacagggcgcgtactatggttgct
 ttgacgcacgtctagaagaaccattattatcatgacattaacctataaaaaataggcgatca
 cgaggccctttcgtcttcaagcagatctgaaaaaaagccgcctcattaggcgggctcagat
 ctgctcatgtttgacagcttatcatcgatgtcgacggtaaccgaattcctcgagctctagaag
 cttgagctcggatccgaattctgaaatccttccctcgatcccgagggttggttgattgttat

FIG. 6A

15/22

tgttgttgttgttcgagctcgaattagtctgcgcgtctttcagggcttcacgacagtctga
cgaccgctggcggttgatcacgcgtagcacggcataaccagaaagcgacatctgcgg
gatgttcggcatgatttcacctttctgggcgttttccatagtggcggaacacgtggatctt
tcgccaaactcttcctogtaagacttcagcgctacggcacccagcggtttgtctttattaacc
gcttcagaccttcacagtcagcagatagttttcagggaactcttttgccagctctttgtt
cggactggcggttaataacctgcgctcagcacgccaaacgaacgggtttggatgggtgacct
tgaaggctcggcagtagcgttacaccataattcactttgctggtgctgatgttggaccatgcc
cacgggcggttgatggctatogctgtttcgcctttattaaaggcagctctcgcgatggagta
atcgggtgtctgcattcatgtgtttgttttaatacaggtcaaccaggaaggctcagaccgcct
tcgcgccagcgttatccacgcccacgtctttaatgtcgtacttgccgttttcatacttgaac
gcataaaccocgtagcagcaatcagcgccagggtgaagtacgggtcttgcagggtgaacat
cagcgcgctcttacctttcgccttcagttctttatccagcgcgggatctcttccaggttt
ttggcggttcggcagcagatctttgttataaatcagcgataacgcttcaacagcgatcggg
taagcaatcagcttgccgttgtaacgtacggcatcccaggtaaacggatacagcttgtctg
gaacgctttgtccgggtgatttcagccaacaggccagattgagcgtagccaccaaacgggt
cgtgtgcccagaagataatgtcaggccatcgccagttgcccaacctgtgggaatttctct
tcagtttatccggatgctcaacggtgactttaattccgggtatctttctcgaatttcttacc
gacttcagcgagaccgttatagcctttatcgcggttaatccagattaccagtttacctctt
cgattttcatatgacctcctaagcatcgatagatcctgtttcctgtgtgaaattgttatccg
ctcacaaattccacacattatacagagccgatgattaattgtcaacagggggatggggagtaag
ctgatcctgtttcctgtgtgaaattgttatccgctcacaattccacacattatacagagccga
tgattaattgtcaacagggggatggggagtaagctcatcgatggatcgatcctgtttcctgt
gtgaaattgttatccgctcacaattccacacattatacagagccggaagcataaaagtgtaaag
cctgggggtgcctaattgagtgagctaaacttacattaattgcgttgcgctcactgcccgcttc
cagtcgggaaacctgtcgtgccaggacaccatcgaattgggtgaaaaacctttcgcgggtatggc
atgatagcgccggaagagagtaaatcagggtgggtgaattgtgaaaccagtaacgtttatag
atgtcgcagagtagtgccggtgtctcttatcagaccgtttcccgcggttgtaaccaggccagc
cacgtttctgcgaaaaacgggaaaaagtgggaagcgcgatggcgagctgaattacattcc
caaccgcgtggcacaacaactggcgggcaaacagtcgttgctgattggcgttgccacctcca
gtctggccctgcacgcgcgtcgcaaatgtcgcggcgattaaatctcgcgccgatcaactg
gggtgccagcgtgggtgtcgatggtagaacgaagcggcgtcgaagcctgtaaagcggcggt

FIG. 6B

16/22

gcacaatcttctcgcgcgaacgcgctcagtgaggctgatcattaactatccgctggatgaccagg
atgccattgctgtggaagctgcctgcactaatgttccggcggtattttcttgatgtctctgac
cagacacccatcaacagttatttttctcccatgaagacggtacgcgactgggctggagca
tctggtcgcattgggtcaccagcaaatcgcgctgttagcgggccattaaagttctgtctcgg
cgcgtctgcgtctggctggctggcataaatatctcactcgaatcaaattcagcogatagcg
gaacgggaaggcgactggagtgccatgtccggttttcaacaacatgcaaattgctgaatga
gggcatcgttcccaactgcgatgctggttgccaacgatcagatggcgctgggcgcaatgcgcg
ccattaccgagtcocggctgcgcgttggtgcggatatctcggtagtgggatacgcacgatacc
gaagacagctcatgttatatcccgccgttaaccacccatcaaacaggattttcgcctgctggg
gcaaacacagcgtggaccgcttgctgcaactctctcagggccaggcggtgaaggcgcaatcagc
tgttgcgcgtctcactgggtgaaaagaaaaaccocctggcgcccaatacgcacaaaccgcctct
ccccgcgcgttgccgattcattaatgcagctggcacgcaggtttccgcactggaaagcgg
gcagtgagcgcacgcacaaatgaagtttagctcactcattagcaccocagcgtttacac
tttatgcttccgcgtctgatggcgtttcggctgatgcaggtgaaaacctctgacacatgcagc
tcccgagacggtcacagcttgctgtgaagcggatgcggggagcagcaagcccgctcagggc
gcgtcagcgggtgttgccgggtgtcggggcgcagccatgaccagtcacgtagcgtatgcg
agtgatactggcttaactatgcggcatcagagcagattgtactgagagtgacacattatgc
ggtgtgaaataccgcacagatgcgtaaggagaaaaataccgcacacaggcgtctctccgcttcc
tcgctcactgactcgtgcgcctcggctcgttcggctgcggcgagcggatcagctcactcaaa
ggcggtaatacgggttatccacagaatcaggggataacgcagggaagaacatgtgagcaaaag
gccagcaaaaggccagggaacgcgtaaaaaggccgcgttgctggcggtttttccataggctccgc
ccccctgacgagcatcacaataatgcagcgtcaagtcagaggtggcgaaaccgcagcagcact
ataagataccaggcgtttccccctggaagctccctcgtgcgcctctcctggttcgcacccctgc
cgcttacgggatacctgtccgcctttctcccttcgggaagcgtggcgctttctcatagctca
cgctgtagggtatctcagttcgggtgtaggtcgttcgctccaagctgggctgtgtgcacgaacc
ccccgttcagcccgaccgctgcgccttatccggtaactatcgtcttgagtcacacccggta
gacacgacttatcgccactggcagcagccactggttaacaggattagcagagcagaggtatgta
ggcgggtgtacagagttcttgaagtggtggcctaactacggctacactagaaggacagttat
tggtatctgcgcctctgctgaagccagttaccttcgggaaaaagagttggtagctcttgatccg
gcaaacacacccgcgtggtagcgggtggttttttgggttgcaagcagcagattacgcgcgaga
aaaaaaggatctcaagaagatcctttgatcttttctacgggggtcgcagctcagtggaacga




FIG. 6C

17/22

aaactcacgttaagggatttttggtcatgagattatcaaaaaggatcttcacctagatccttt
taaattaaaaatgaagttttaaatcaatctaaagtatatatgagtaaacttggtctgacagt
taccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcggttcattccatagt
tgctgactccccgtcgtgtagataactacgatacgggaggggttaccatctggccccagtg
ctgcaatgataaccgcgagacccacgctcacggctccagatttatcagcaataaaccagcca
gcgggaagggccgagcgcagaagtggtcctgcaactttatccgcctccatccagctctattaa
ttggttgcgggaagctagagtaagtagttcgccagttaatagtttgcgcaacgttggtgcca
ttgctgcag

FIG. 6D

18/22



19/22




Fig. 7D




Fig. 7E

20/22

Fig. 7F





Fig. 7G



21/22




Fig. 8A




Fig. 8B



Fig. 8C

22/22

M 1 2 3 4 5 6 7 8 9 10 11 12 13





Fig. 8D

Fig. 8E



SEQUENCE LISTING

<110> Neose Technologies, Inc.
 Johnson, Karl
 Bezila, Dan
 Ngo, Winnie
 Hakes, David

<120> Vectors for Recombinant Protein Expression in E. coli

<130> 040853-01-5090WO

<140> Not Yet Assigned

<141> 2005-01-06

<150> U.S. 60/535,263

<151> 2004-01-09

<160> 13

<170> PatentIn version 3.2

<210> 1

<211> 5039

<212> DNA

<213> Artificial

<220>

<223> Custom DNA vector

<400> 1

gcacgtggtg gtcacgtctg tcgtttggta tggcttcatt cagctccggt tcccaacgat	60
caaggcgagt tacatgatcc cccatgttgt gcaaaaaagc ggtagctcc ttcggtcctc	120
cgatcggggg gggggggaaa gccacgttgt gtctcaaaat ctctgatgtt acattgcaca	180
agataaaat atatcatcat gaacaataaa actgtctgct tacataaaca gtaatacaag	240
gggtgttatg agccatattc aacgggaaac gtcttgcctc aggcgcgat taaattccaa	300
catggatgct gatttatatg ggtataaatg ggctcgcgat aatgtcgggc aatcaggtgc	360
gacaatctat cgactgtatg ggaagccga tgcgccagag ttgtttctga aacatgcaa	420
aggtagcggt gccaatgatg ttacagatga gatggtcaga ctaaactggc tgacgggaatt	480
tatgcctctt cggaccatca agcattttat ccgtactcct gatgatgcat gggtactcac	540
cactgcgata cccgggaaaa cagcattcca ggtattagaa gaatatcctg attcaggtga	600
aaatattgtt gatgcgctgg cagtgttcct gcgccggttg cattcgatcc ctgtttgtaa	660
ttgtcctttt aacacgcgac gcgtatttcg tctcgtcag gcgcaatcac gaatgaataa	720
cggtttggtt gatgcgagtg attttgatga cgagcgtaat ggctggcctg ttgaacaagt	780
ctggaagaa atgcataagc tattgccatt ctcacccgat tcagtcgtca ctcatgggtga	840
tttctcactt gataacctta tttttgacga ggggaaatta ataggttgta ttgatgttgg	900

acgagtcgga atcgcagacc gataccagga tcttgccatc ctatggaact gcctcgggtga	960
gttttctcct tcattacaga aacggctttt tcaaaaatat ggtattgata atcctgatat	1020
gaataaattg cagtttcatt tgatgctcga tgagttttcc taaagtacta ctcttccttt	1080
ttcaatatta ttgaagcatt tatcagggtt attgtctcat gagcggatgc atatttgaat	1140
gtatttagaa aaataaacaa ataggggttc cgcgcacatt tccccgaaaa gtgccacctg	1200
acgatgaaat tgtaaacggt aatattttgt taaaattcgc gttaaatttt tgttaaatca	1260
gctcattttt taaccaatag gccgaaatcg gcaaaatccc ttataaatca aaagaatagc	1320
ccgagatagg gttgagtgtt gttccagttt ggaacaagag tccactatta aagaacgtgg	1380
actccaacgt caaagggcga aaaaccgtct atcaggggcga tggcccacta cgtgaaccat	1440
cacccaaatc aagttttttg gggctcgaggt gccgtaaaag tctaaatcgg aacctaaag	1500
ggagcccccg atttagagct tgacggggaa agccggcgaa cgtggcgaga aagggaaggga	1560
agaaagcgaa aggagcgggc gctagggcgc tggcaagtgt agcggtcacg ctgcgcgtaa	1620
ccaccacacc cgcgcgcgtt aatgcgcgcg tacaggggcg gtactatggt tgctttgacg	1680
catcgtctaa gaaaccatta ttatcatgac attaacctat aaaaatagcg gtatcacgag	1740
gccctttcgt cttcaagcag atctgaaaaa aaagcccgtc cattaggcgg gctcagatct	1800
gctcatgttt gacagcttat catcgatgc gacggtacg aattcctcga gtctagaaag	1860
cttgagctcg gatcccatat gacctctaa gcatcgatgg atcctgttcc ctgtgtgaaa	1920
ttgttatccg ctccacaatt cacacattat acgagccgat gattaattgt caacaggggg	1980
atggggagta agctgatcct gtttcctgtg tgaatttgtt atccgctcac aattccacac	2040
attatacgag ccgatgatta attgtcaaca gggggatggg gagtaagctc atcgatggat	2100
cgatcctggt tctctgtgtg aattgttatt cgctcacaat tccacacatt atacgagccg	2160
gaagcataaa gtgtaaagcc tgggggtgct aatgagttag ctaacttaca ttaattgcgt	2220
tgcgctcact gcccgcttcc cagtcgggaa acctgtcgtg ccaggacacc atcgaatggt	2280
gcaaaacctt tcgcggtatg gcatgatagc gcccggaaga gagtcaattc aggggtgggtga	2340
atgtgaaacc agtaacgltt tacgatgtcg cagagtatgc cgggtgtctc tatcagaccg	2400
tttcccggtt ggtgaaccag gccagccacg tttctgcgaa aacgcgggaa aaagtgggaag	2460
cggcgatggc ggagctgaat tacattccca accgcgtggc acaacaactg gcggggcaaac	2520
agtcgttgct gattggcgtt gccacctcca gtctggccct gcacgcgcg tcgcaaatgt	2580
tcgcgccgat taaatctcgc gccgatcaac tgggtgccag cgtgggtggtg tcgatggtag	2640
aacgaagcgg cgtcgaagcc tgtaaagcgg cgggtgcaca tcttctcgcg caacgcgtca	2700
gtgggctgat cattaactat ccgctggatg accaggatgc cattgtctgtg gaagctgcct	2760

gcactaatgt tccggcggtta tttcttgatg tctctgacca gacaccctac aacagtatta 2820

ttttctccca tgaagacggt acgcgactgg gcgtggagca tctggctgca ttgggtcacc 2880

agcaaatcgc gctgttagcg gggccattaa gttctgtctc ggcgcgtctg cgtctggctg 2940

gctggcataa atatctcact cgcaatcaaa ttcagccgat agcggaacgg gaaggcgact 3000

ggagtgccat gtccggtttt caacaaacca tgcaaatgct gaatgagggc atcgttccca 3060

ctgcgatgct ggttgccaac gatcagatgg cgctgggcgc aatgcgcgcc attaccgagt 3120

ccgggctcgc cgttggtgog gatattctcg tagtgggata cgacgatacc gaagacagct 3180

catgttatat ccgcgcgtta accaccatca aacaggattt tcgcctcgtg gggcaacca 3240

gcgtggacgg cttgctgcaa ctctctcagg gccaggcggg gaagggaacat cagctgttgc 3300

ccgtctcact ggtgaaaaga aaaaccaccc tggcgcccaa tacgcaaacc gcctctcccc 3360

gcgcgttggc cgattcatta atgcagctgg cagcacaggt tccccgactg gaaagcgggc 3420

agtgagcgca acgcaattaa tgtaagttag ctactcatt aggcacccca ggctttacac 3480

tttatgcttc cggctcgtat ggcgtttcgg tgatgacggg gaaaacctct gacacatgca 3540

gtccccggag acggctcacg cttgtctgta agcggatgcc gggagcagac aagcccgta 3600

gggcgcgtca gggggtgttg gggggtgtcg ggggcagacc atgaccaggt cagtagcgca 3660

tagcggagtg tatactggct taactatgcg gcatcagagc agattgtact gagagtgcac 3720

cattatgcgg tgtgaaatc cgacacagat cgtaaggaga aaatacgcga tcaggcgctc 3780

ttccgcttcc tcgctcactg actcgtcgcg ctccgctcgt cggctcggcg gagcggtatc 3840

agctcactca aaggcggtaa tacggttatc cacagaatca ggggataacg caggaaagaa 3900

catgtgagca aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt tgcgtggcgtt 3960

tttccatagg ctccgcccc ctgacgagca tcacaaaaat cgacgctcaa gtcagagggtg 4020

gcgaaacccg acaggactat aaagatacca ggcgtttccc cctggaagct ccctctgctg 4080

ctctcctggt ccgacccctc cgcttacccg atacctgtcc gcctttctcc ctccgggaag 4140

cgtggcgctt tctcatagct cagcgtgtag gtatctcagt tcggtgtagg tcgttcgctc 4200

caagctgggc tgtgtgcacg aacccccgtt tcagcccgac cgctgcgcct tatccggtaa 4260

ctatcgtctt gagtccaacc cggttaagaca cgacttatcg ccactggcag cagccactgg 4320

taacaggatt agcagagcga ggtatgtagg cgggtctaca gagtcttga agtggtggcc 4380

taactacggc tacactagaa ggacagtatt tggatatcgc gctctgctga agccagttac 4440

cttcggaaaa agagtttgta gctcttgatc cggcaaaaa accaccgctg gtacgggtgg 4500

ttttttgtt tgcaagcagc agattacgcg cagaaaaaaa ggatctcaag aagatccttt 4560

gatctttttct acgggggtctg acgctcagtg gaacgaaaac tcacgttaag ggatttttgg	4620
catgagatta tcaaaaagga tcttcaccta gatcctttta aattaaaaat gaagttttta	4680
atcaatctaa agtatatatg agtaaacttg gtctgacagt taccaatgct taatcagtga	4740
ggcaccatc tcagcgatct gtctatttgc ttcattccata gttgcctgac tccccgtcgt	4800
gtagataact acgatacggg agggcttacc atctggcccc agtgctgcaa tgataccgcg	4860
agacccacgc tcaccggctc cagatttacc agcaataaac cagccagccg gaagggccga	4920
gcgcagaagt ggtcctgcaa ctttatccgc ctccatccag tctattaatt gttgccggga	4980
agctagagta agtagttcgc cagttaatag tttgcgcaac gttgttgcca ttgctgcag	5039

<210> 2
 <211> 5039
 <212> DNA
 <213> Artificial

<220>
 <223> Custom DNA vector

<400> 2 gcatcgtggt gtcacgctcg tcgttttgga tggcttcatt cagctccggt toccaacgat	60
caaggcgagt tacatgatcc cccatgttgt gcaaaaaagc ggtagctcc ttcgggtctc	120
cgatcggggg gggggggaaa gccacgttgt gtctcaaat ctctgatgtt acattgcaca	180
agataaaaat atatcatcat gaacaataaa actgtctgct tacataaaca gtaatacaag	240
gggtgttatg agccatattc aacgggaaac gtcttgctcc aggcgcgat taaattccaa	300
catggatgct gatttatatg ggtataaatg ggctcgcgat aatgcgggc aatcaggtgc	360
gacaatctat cgactgtatg ggaagccga tgcgcagag ttgtttctga aacatggcaa	420
aggtagcggt gccaatgatg ttacagatga gatggtcaga ctaaactggc tgacgggaatt	480
tatgcctctt ccgaccatca agcattttat ccgtactcct gatgatgcatt ggttactcac	540
cactgcgata cccgggaaaa cagcattcca ggtattagaa gaatatcctg attcaggtga	600
aaatatgtt gatgcgctgg cagtgttcct gcgccggttg cattcgattc ctgtttgtaa	660
ttgtcctttt aacagcgatc gcgtatttgc tctcgtcag gcgcaatcac gaatgaataa	720
cggtttggtt gatgcgagtg attttgatga cgagcgtaat ggctggcctg ttgaacaagt	780
ctggaaagaa atgcataagc tattgccatt ctacccgat tcagtcgtca ctcatgggtga	840
tttctcattt gataacctta tttttgacga ggggaaatta ataggttgta ttgatgttgg	900
acgagtcgga atcgcagacc gataccagga tcttgccatc ctatggaact gcctcgggtga	960
gttttctcct tcattacaga aacggctttt tcaaaaatat ggtattgata atcctgatat	1020
gaataaatgt cagtttcatt tgatgctoga tgagtttttc taaagtacta ctcttctctt	1080

ttcaatatta ttgaagcatt tatcagggtt attgtctcat gagcggatag atatttgaat 1140
 gtatttagaa aataaaca ataggggttc cgcgcacatt tccccgaaa gtgccacctg 1200
 acgatgaat tgtaaacgtt aatattttgt taaaattcgc gttaaatatt tgtaaataca 1260
 gctcattttt taaccaatag gccgaaatcg gcaaaatccc ttataaatca aagaatatagc 1320
 ccgagatagg gttgagtgtt gttccagttt ggaacaagag tccactatta aagaacgtgg 1380
 actccaacgt caaagggcga aaaacggtct atcagggcga tggccacta cgtgaaccat 1440
 caccocaaatc aagttttttg gggctcgaggt gccgtaaagc tctaaatcgg aacctaaag 1500
 ggagcccccg atttagagct tgacggggaa agccggcgaa cgtggcgaga aaggaaggga 1560
 agaaagcgaa aggagcgggc gctagggcgc tggcaagtgt agcggtcacg ctgcgcgtaa 1620
 ccaccacacc cgcgcgcgtt aatgcgcgcg tacagggcgc gtactatggt tgctttgacg 1680
 catcgtctaa gaaaccatta ttatcatgac attaacctat aaaaatagcg gtatcacgag 1740
 gccctttcgt cttcaagcag atctgaaaaa aaagcccgct cattaggcgg gctcagatct 1800
 gctcatgttt gacagcttat catcgatgac gacggtagcg aattcctcga gtctagaag 1860
 cttgagctcg gatcccatat gacctctaa gcatcgatag atcctgttct ctgtgtgaaa 1920
 ttgttatccg ctacacattc cacacattat acgagccgat gattaattgt caacaggggg 1980
 atggggagta agctgatctt gtttctgtg tgaatttgtt atccgctcac aattccacac 2040
 attatacgag ccgatgatta attgtcaaca gggggatggg gagtaagctc atcgatggat 2100
 cgatcctggt tctctgttga aattgttctc cgctcacaat tccacacatt atacgagccg 2160
 gaagcataaa gtgtaaaagc tggggtgcct aatgagtgag ctaacttaca ttaattgcgt 2220
 tgcgctcact gcccgcttct cagtcgggaa acctgtcgtg ccaggacacc atcgaatggt 2280
 gcaaaacctt tcgcggtatg gcatgatagc gcccggaaga gagtcaattc aggggtggtga 2340
 atgtgaaacc agtaacgtta tacgatgtcg cagagtatgc cgggtgtctct tatcagaccg 2400
 tttcccgctt ggtgaaccag gccagccacg tttctgcgaa aacgcgggaa aaagtgggaag 2460
 cggcgatggc ggagctgaat tacattccca accgcgtggc acaacaactg gcgggcaaac 2520
 agtcgttctt gattggcgtt gccacctcca gtctggccct gcacgcgcgc tcgcaaatgt 2580
 tcgcggcgat taaatctcgc gccgatcaac tgggtgccag cgtgggtggtg tcgatggtag 2640
 aacgaagcgg cgtcgaagcc tgtaaaagcg cgggtgcaca tcttctcgcg caacgcgtca 2700
 gtgggctgat cattaactat ccgctggatg accaggatgc cattgctgtg gaagctgcct 2760
 gcactaatgt tccggcgcta tttcttgatg tctctgacca gacaccatc aacagtatta 2820
 ttttctccca tgaagacggt acgcgactgg gcgtggagca tctggtcgca ttgggtcacc 2880

agcaaatcgc gctgttagcg ggcccattaa gttctgtctc ggcgcgtctg cgtctggtcg	2940
gctggcataa atatctcact cgcaatcaaa ttcagccgat agcgggaacgg gaaggcgact	3000
ggagtgccat gtccgggtttt caacaaacca tgcaaatgct gaatgagggc atcgtttccca	3060
ctgcgatgct ggttgccaac gatcagatgg cgctggggcg aatgcgcgcc attaccgagt	3120
cggggctcgc cgttggttgcg gatatactcg tagtgggata cgacgatacc gaagacagct	3180
catgttatat ccgcgcgtta accaccatca aacaggattt tcgcctgctg gggcaaacca	3240
gcgtggacgg cttgtctgca ctctctcagg gccaggcggg gaagggcaat cagctgttgc	3300
ccgtctcact ggtgaaaaga aaaaccaccc tggcgcccaa tacgcaaac gccctctccc	3360
gcgcgttggc cgattcatta atgcagctgg cagcagcagg tcccgcactg gaaagcgggc	3420
agtgagcgca acgcaattaa tgtaagttag ctactcatt aggcacccca ggctttacac	3480
tttatgcttc cggctcgtat ggcgtttcgg tgatgacggg gaaaacctct gacacatgca	3540
gtccccggag acggtcacag cttgtctgta agcggatgcc gggagcagac aagcccgta	3600
gggcgcgtca gcgggtgttg gcgggtgtcg gggcgagacc atgaccagc cagtagcgga	3660
tagcggagtg tatactggct taactatcgc gcacagagc agattgtact gagagtgcac	3720
cattatgcgg tgtgaaatac cgcacagatg cgtaaggaga aaataccgca tcaggcgctc	3780
ttccgcttc tcgctcactg actcgtcgcg ctcggtcgtt cggctgcggc gagcggtatc	3840
agctcactca aaggcggtaa tacggttacc cacagatca ggggataacg caggaaagaa	3900
catgtgagca aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt tgctggcgct	3960
tttccatagg ctccgcccc ctgacgagca tcacaaaaat cgacgcctca gtcagaggtg	4020
gcgaaacccg acaggactat aaagatacca ggcgtttccc cctggaagct ccctcgtgcg	4080
ctctctgctt ccgacctcgc cgcttacggg atacctgtcc gcctttctcc ctccgggaag	4140
cgtggcgctt tctcatagct cagcctgtag gtatctcagt tcggtgtagg tcgttcgctc	4200
caagctgggc tgtgtgcacg aacccccgt tcagcccgac cgctgcgcct tatccggtaa	4260
ctatcgtctt gagtccaacc cggtaagaca cgacttatcg ccactggcag cagccactgg	4320
taacaggatt agcagagcga ggtatgtagg cggtgctaca gattcttga agtggtggcc	4380
taactacggc tacactagaa ggacagtatt tggatatcgc gctctgtga agccagttac	4440
cttcggaaaa agagttggta gctcttgatc cggcaaaaa accaccgctg gtacgggtgg	4500
tttttttgtt tgcaagcagc agattacgcg cagaaaaaaa ggtatctcaag aagatccttt	4560
gatcttttct acggggtctg acgctcagtg gaacgaaaa tcacgtaag ggatttttgt	4620
catgagatta tcaaaaagga tcttcaccta gatccttta aattaaaaat gaagttttta	4680
atcaatctaa agtatatatg agtaaacctg gtctgacagt taccaatgct taatcagtga	4740

ggcacctatc	tcagcgatct	gtctatttcg	ttcatccata	gttgccctgac	tccccgtcgt	4800
gtagataact	acgatacggg	agggccttacc	atctggcccc	agtgcctgcaa	tgataccgcg	4860
agacccacgc	tcacgggctc	cagattttatc	agcaataaac	cagccagccg	gaagggccga	4920
gcgcagaagt	ggctcctgcaa	ctttatccgc	ctccatccag	tctattaatt	gttgccggga	4980
agctagagta	agtagttcgc	cagttaatag	tttgcgcaac	gttggttgcca	ttgctgcag	5039

<210> 3
 <211> 6209
 <212> DNA
 <213> Artificial

<220>
 <223> Custom DNA vector

<400> 3	
gcacgtgggt	gtcacgctcg
tcgttttggt	tggcttcatt
cagctccggt	tcccaacgat
60	
caaggcgagt	tacatgatcc
cccatgttgt	gcaaaaaagc
ggttagctcc	ttcgggtcctc
120	
cgatcggggg	gggggggaaa
gccacgttgt	gtctcaaaat
ctctgatgtt	acattgcaca
180	
agataaaat	atatcatcat
gaacaataaa	actgtctgct
tacataaaca	gtaatacaag
240	
gggtgttatg	agccatatct
aacgggaaac	gtcttgctcc
aggccgcgat	taaattccaa
300	
catggatgct	gatttatatg
ggtataaatg	ggctcgcgat
aatgtcgggc	aatcaggtgc
360	
gacaatctat	cgactgtatg
ggaagcccca	tgcgccagag
ttgtttctga	aacatggcaa
420	
aggtagcggt	gccaatgatg
ttacagatga	gatggtcaga
ctaaactggc	tgacggaatt
480	
tatgcctctt	ccgaccatca
agcattttat	ccgtactcct
gatgatgcat	ggttactcac
540	
caactgogac	cccgggaaaa
cagcattcca	ggtattagaa
gaatatcctg	attcaggtga
600	
aaatattgtt	gatgcgctgg
cagtgttcct	gcgccgggtg
cattcgatcc	ctgtttgtaa
660	
ttgtcctttt	aacagcgatc
gcgtatttcg	tctcgctcag
gcgcaatcac	gaatgaataa
720	
cggtttggtt	gatgcgagtg
attttgatga	cgagcgtaat
ggctggcctg	ttgaacaagt
780	
ctggaaagaa	atgcataagc
tattgccatt	ctcaccggat
tcagtcgtca	ctcatggtga
840	
ttttctcact	gataacctta
tttttgacga	ggggaaatta
ataggttgta	ttgatgttgg
900	
acgagtcgga	atcgcagacc
gataccagga	tcttgccatc
ctatggaact	gcctcggtga
960	
gttttctcct	tcattacaga
aacggccttt	tcaaaaaaat
ggtattgata	atcctgatata
1020	
gaataaatgg	cagtttcatt
tgatgctcga	tgagtttttc
taaaagta	ctcttccttt
1080	
ttcaatatta	ttgaagcatt
tatcagggtt	attgtctcat
gagcggatac	atatttgaat
1140	
gtatttagaa	aaataaacaa
ataggggttc	cgcgcacatt
tccccgaaaa	gtgccacctg
1200	

acgatgaaat	tgtaaacggt	aatatatttgt	taaaattcgc	gttaaatttt	tgtaaataca	1260
gctcattttt	taaccaatag	gccgaaatcg	gcaaaatccc	ttataaatca	aaagaatagc	1320
ccgagatagg	gttgagtgtt	gttcacgttt	ggaacaagag	tccactatta	aagaacgtgg	1380
actccaacgt	caaaggcgga	aaaacggtct	atcagggcga	tggcccacta	cgtgaaccat	1440
cacccaaatc	aagttttttt	gggtcgaggt	gccgtaagc	tctaaatcgg	aaccctaaag	1500
ggagcccccg	atttagagct	tgacggggaa	agccggcgaa	cgtggcgaga	aaggaaggga	1560
agaaagcgaa	aggagcgggc	gctagggcgc	tggcaagtgt	agcggtcacg	ctgcgcgtaa	1620
ccaccacacc	cgcgcgctt	aatgcgccgc	tacagggcgc	gtactatggt	tgctttgacg	1680
catcgcttaa	gaaaccatta	ttatcatgac	attaacctat	aaaaatagcg	gtatcacgag	1740
gccctttcgt	cttcaagcag	atctgaaaaa	aaagcccgct	cattaggcgg	gctcagatct	1800
gctcatgttt	gacagcttat	catcgatgtc	gacggtaccg	aattcctcga	gtctagaag	1860
cttgagctcg	gatccgaatt	ctgaaatcct	tcctcgatc	ccgaggttgt	tgttattgtt	1920
attgttgttg	ttgttcgagc	tcgaattagt	ctgcgcgtct	ttcagggtct	catcgacagt	1980
ctgacgaccg	ctggcggcgt	tgataccgc	agtacgcacg	gcataaccaga	aagcggacat	2040
ctgcccgatg	ttcgccatga	tttcaccttt	ctggcggttt	tccatagtgg	cggcaatacg	2100
tggatctttc	gccaaactct	cctcgtaaga	cttcagcgct	acggcaccca	gcggtttgtc	2160
tttattaacc	gcttcacagc	cttcacagct	cagcagatag	ttttcgagga	actctttttg	2220
cagctctttg	ttcggaactg	cggcgtaaat	acctgcgctc	agcacgccea	cgaacggttt	2280
ggatggttga	cccttgaagg	tcggcagtac	cgttacacca	taattcactt	tgctggtgtc	2340
gatgttgga	catgcccacg	ggccgttgat	ggctcatcgct	gtttcgcttt	tattaaaggc	2400
agcttctcgg	atggagtaat	cgggtgtctgc	attcatgtgt	ttgtttttaa	tcagggtcaac	2460
caggaaggtc	agaccgcgtt	tcgcgccacg	gttatccacg	cccacgtctt	taatgtcgta	2520
cttgccgttt	tcataactga	acgcataacc	cccgtcagca	gcaatcagcg	gccagggtgaa	2580
gtacggttct	tgacggttga	acatcagcgc	gctcttacct	ttcgctttca	gttctttatc	2640
cagcgcgggg	atctcttccc	aggttttttg	cgggttcggc	agcagatctt	tggtataaat	2700
cagcgataac	gcttcaacag	cgatcgggta	agcaatcagc	ttgccgttgt	aacgtacggc	2760
atcccaggta	aacggatata	gcttgtcctg	gaacgctttg	tccggggtga	tttcagccaa	2820
caggccagat	tgagcgtagc	caccaaagcg	gtcgtgtgcc	cagaagataa	tgtcaggggc	2880
atcgccagtt	gccgcaacct	gtgggaattt	ctcttcaggt	ttatccggat	gctcaacggt	2940
gactttaatt	ccggtatctt	tctcgaattt	cttaccgact	tcagcgagac	cggttatagcc	3000
tttatcgccg	ttaatccaga	ttaccagttt	accttctctg	attttcatat	gacctcctaa	3060

gcatcgatag atcctgtttc ctgtgtgaaa ttgttatccg ctacacaattc cacacattat 3120
 acgagccgat gattaattgt caacaggggg atggggagta agctgatcct gtttccctgtg 3180
 tgaattgtt atccgctcac aattccacac attatacgag ccgatgatta attgtcaaca 3240
 ggggatggg gagtaagctc atcgatggat cgalcctgtt tcctgtgtga aattgttattc 3300
 cgctcacaat tccacacatt atacgagccg gaagcataaa gtgtaaagcc tgggggtccct 3360
 aatgagttag ctaacttaca ttaattgctg tgcgtcact gccgccttc cagtcgggaa 3420
 acctgtctg ccaggacacc atcgaatggt gcaaacctt tcgcggtatg gcgatgatgc 3480
 gcccggaaga gagtcaattc aggggtgtga atgtgaaacc agtaacgtta tacgatgtcg 3540
 cagagtatgc cgtgtctct tatcagaccg ttcccgctg ggtgaaccag gccagccacg 3600
 tttctgcgaa aacgcgggaa aaagtggaag cggcgatggc ggagctgaat tacattccca 3660
 accgcgtggc acaacaactg gcgggcaaac agtcgttctg gattggcgtt gccacctcca 3720
 gtctggccct gcacgcgcg tcgcaaattg tcgcggcgat taaatctcgc gccgatcaac 3780
 tgggtgccag cgtgtgtgtg tcgatggtag aacgaagcgg cgtcgaagcc tgtaaagcgg 3840
 cggtgccaaa tcttctcgcg caacgcgtca gtgggctgat cattaactat ccgctggatg 3900
 accaggatgc cattgtctg gaagctgcct gcactaatgt tccggcggtta tttcttgatg 3960
 tctctgacca gacaccatc aacagtatta tttctccca tgaagacggt acgcgactgg 4020
 gcgtggagca tctggctgca ttgggtcacc agcaaatcgc gctgttagcg gggccattaa 4080
 gttctgtctc ggcgcgtctg cgtctggctg gctggcataa atatctcact cgcaatcaaa 4140
 ttcagccgat agcgaacggg gaaggcgact ggagtgccat gtccggtttt caacaaacca 4200
 tgcaaatgct gaatgagggc atcgttccca ctgcgatgct ggttgccaac gatcagatgg 4260
 cgctgggcgc aatgcgcgcc attaccgagt ccgggctcgc cgttggtgcg gatattctcg 4320
 tagtgggata cgacgatacc gaagacagct catgttatat ccgcgcgtta accaccatca 4380
 aacaggattt tcgcctctg gggcaaacca gcgtggacgg cttgctgcaa ctctctcagg 4440
 gccaggcggg gaagggcaat cagctgttgc ccgtctcact ggtgaaagaa aaaaccaccc 4500
 tgggcccaca tacgcaaacc gcctctcccc gcgcgttggc cgattcatta atgcagctgg 4560
 cacgacaggt ttccgactg gaaagcgggc agtgagcgca acgcaattaa tgtaagtttag 4620
 ctactcatt aggcacccca ggctttacac ttatgtcttc cgctcgtat ggcgttttcgg 4680
 tgatgacggt gaaaacctct gacacatgca gctcccgagg acggtcacag cttgtctgta 4740
 agcggatgcc gggagcagac aagcccgta gggcgcgtca gcgggtgttg gcgggtgtcg 4800
 gggcgacgcc atgaccagc cactagcga tagcggagtg tatactggct taactatcgg 4860

gcatcagagc agattgtact gagagtgcac cattatgcgg tgtgaaatac cgcacagatg 4920
 cgttaaggaga aaataaccga tcaggcgctc ttccgcttcc tcgctcactg actcgtgctg 4980
 ctccgtcggtt cggctgcggc gagcggtatc agctcactca aaggcggtaa tacggttatc 5040
 cacagaatca ggggataacg caggaaagaa catgtgagca aaagggccagc aaaagggccag 5100
 gaaccgtaaa aaggccgcgt tgctggcggtt ttcccatagg ctccgcccc ctgacgagca 5160
 tcacaaaaat cgacgctcaa gtcagagggtg gcgaaacccg acaggactat aaagatacca 5220
 ggcgtttccc cctggaagct ccctcgctgc ctctcctgtt ccgacctgc cgcttaccgg 5280
 atacctgtcc gcctttctcc ctccgggaag cgtggcgctt tctcatagct cacgctgtag 5340
 gtatctcagt tcgggtgagg tcgttcgctc caagctgggc tgtgtgcacg aacccccctg 5400
 tcagcccgac cgctgcgcct tatccggtaa ctatcgtctt gagtccaacc cggtaagaca 5460
 cgacttatcg ccactggcag cagccactgg taacaggatt agcagagcga ggtatgtagg 5520
 cgggtctaca gagttcttga agtgggtggc taactacggc tacactagaa ggacagtatt 5580
 tggatatctg cctctgtcta agccagttac ctccgaaaa agagttggta gctcttgatc 5640
 cggcaaacaa accaccctg gtagcgggtg tttttttgtt tgcaagcagc agattacgctg 5700
 cagaaaaaaaa ggaatcctaa agatccctt gatcttttct acggggctctg acgctcagtg 5760
 gaacgaaaac tcacgttaag ggatttttgt catgagatta tcaaaaagga tcttcaccta 5820
 gatcctttta aattaaaaat gaagttttaa atcaatctaa agtatatatg agtaaaacttg 5880
 gtctgacagt taccaatgct taatcagtg ggcacctatc tcagcgatct gtctatttcg 5940
 ttcatccata gttgcctgac tccccgtcgt gtagataact acgatacggg agggcttacc 6000
 atctggcccc agtgctgcaa tgataccgcg agaccacgc tcaccggctc cagatttatc 6060
 agcaataaac cagccagcgc gaagggccga gcgcagaagt ggtcctgcaa ctttatccgc 6120
 ctccatccag tctattaatt gttgcggga agctagagta agtagttcgc cagttaatag 6180
 tttagcgaac gttgttgcca ttgctgcag 6209

<210> 4
 <211> 29
 <212> DNA
 <213> Artificial

<220>
 <223> 5' modified restriction site

<400> 4
 attccaattc gatcgggggg ggggggaaa

29

<210> 5
 <211> 31

<212> DNA
 <213> Artificial

 <220>
 <223> 3' modified restriction site

 <400> 5
 attccaagta gtacttttaga aaaactcattc g 31

 <210> 6
 <211> 83
 <212> DNA
 <213> Artificial

 <220>
 <223> 5' modified multiple cloning site

 <400> 6
 atcgatcgac atatgggagc cgagctcaag ctttctagac tcgaggaatt cggtaccgctc 60
 gacatcgatg ataagctgac aaa 83

 <210> 7
 <211> 31
 <212> DNA
 <213> Artificial

 <220>
 <223> 3' modified multiple cloning site

 <400> 7
 attccaagta gtactactct tcctttttca a 31

 <210> 8
 <211> 32
 <212> DNA
 <213> Artificial

 <220>
 <223> 5' PCR primer for pcWIN2 construct

 <400> 8
 caattatata gatctatcga tgcttaggag gt 32

 <210> 9
 <211> 36
 <212> DNA
 <213> Artificial

 <220>
 <223> 3' PCR primer for pcWIN2 construct

 <400> 9
 ttgccttatt ctagatcatt agtgggtgatg gtgggtg 36

 <210> 10

<211> 6659
 <212> DNA
 <213> Artificial

<220>
 <223> Custom DNA vector pMS39

```

<400> 10
tcgccttccc gttccgctat cggctgaatt tgattgcgag tgagatattt atgccagcca      60
gccagacgca gacgcgccga gacagaactt aatgggcccc ctaacagcgc gatttgcttg      120
tgacccaatg cgaccagatg ctccacgccc agtcgcgtac cgtttcatg ggagaaaaata      180
atactgttga tgggtgtctg gtcagagaca tcaagaaata acgcgcgaac attagtgcag      240
gcagcttcca cagcaatggc atcctggcca tccagcggat agttaatgat cagcccactg      300
acgcgttgcg cgagaagatt gtgcaccgcc gctttacagg cttcgacgcc gcttcgttct      360
accatcgaca ccaccacgct ggcaccagtg tgatcggcgc gagatttaat cgccgcgaca      420
atttgcgacg gcgcgtgcag ggcacagactg gaggtggcaa gcgcaatcag caacgactgt      480
ttgcccgcca gttgttgtgc cagcgggttg ggaatgtaat tcagctccgc catcgccgct      540
tccacttttt cccgcgtttt cgcagaaaag tggctggcct ggttcaccac gcgggaaacg      600
gtctgataag agacaccggc atactctgcg acatcgtata acgttactgg ttteacattc      660
accaccctga attgactctc ttccgggcgc tatcatgccca taccgcgaaa ggttttgcac      720
cattcgatgg tgtcctggca cgacaggttt cccgactgga aagcgggcag tgagcgaac      780
gcaattaatg taagttagct cactcattag gcaccccagg ctttacactt tatgcttccg      840
gctcgtataa tgtgtggaat tgtgagcgga taacaatttc acacaggaaa caggatcgat      900
ccatcgatga gcttactccc catccccctg ttgacaatta atcatcggtc cgtataatgt      960
gtggaattgt gagcgggataa caatttcaca caggaaacag gatcagctta ctcccatcc      1020
ccctgttgac aattaatcat cggctcgtat aatgtgtgga attgtgagcg gataacaatt      1080
tcacacagga aacaggatct atcgatgctt aggaggtcat atgaaaatcg aagaaggtaa      1140
actggtaatc tggattaacg gcgataaagg ctataacggg ctcgctgaag tcggtaaaga      1200
attcgagaaa gataccggaa ttaaagtcac cgttgagcat ccggataaac tggaaagaaa      1260
attcccacag gttgcggcaa ctggcgatgg ccctgacatt atcttctggg cacacgaccg      1320
ctttggtggc tacgctcaat ctggcctgtt ggctgaaatc accccggaca aagcgttcca      1380
ggacaagctg tatccgttta cctgggatgc cgtacgttac aacggcaagc tgattgctta      1440
cccgatcgct gttgaagcgt tatcgctgat ttataacaaa gatctgctgc cgaaccgcgc      1500
aaaaacctgg gaagagatcc cggcgctgga taaagaactg aaagcgaagc gtaagagcgc      1560
gctgatgttc aacctgcaag aaccgtactt cacctggcgc ctgattgctg ctgacggggg      1620

```

ttatgcgttc	aagtatgaaa	acggcaagta	cgacattaaa	gacgtgggcg	tggataacgc	1680
tggcgcgaaa	gcgggtctga	ccttcctcgt	tgacctgatt	aaaaacaaac	acatgaatgc	1740
agacaccgat	tactccatcg	cagaagctgc	ctttaataaa	ggcgaaacag	cgatgaccat	1800
caacggcccg	tgggcatggt	cacacatcga	caccagcaaa	gtgaattatg	gtgtaacggc	1860
actgcgcacc	ttcaagggtc	aacctcccaa	accgttcgtt	ggcgtgctga	gcgcaggat	1920
taacgccgcc	agtcggaaca	aagagctggc	aaaagagtgc	ctcgaaaact	atctgctgac	1980
tgatgaagg	ctggaagcgg	ttaataaaga	caaaccgctg	ggcgccgtag	cgctgaagtc	2040
ttacgaggaa	gagttggcga	aagatccacg	tattgccgcc	actatggaaa	acgcccgaaa	2100
agggtgaaat	atgccgaaca	tccgcagat	gtccgctttc	tgggtatgcc	tgcgtaactgc	2160
ggatgatcaac	gccgccagcg	gtcgtcagac	tgctgatgaa	gccctgaaag	acgcgcagac	2220
taattcgagc	tcgaacaaca	acaacaataa	caataacaac	aacctcgggg	tcgaggggaag	2280
gatttcagaa	ttcggatcta	ttgtggcgac	cggcggcacc	accaccacgc	cgaccgccgc	2340
cggctccggc	agcgtgacct	cgaccagcaa	aaccaccgcg	accgcgagca	aaaccagcac	2400
cagcacctca	tcaacctcct	gtaccacccc	gaccgcgggtg	cggtgacct	tcgatctgac	2460
cgcgaccacc	acctacggcg	aaaacatcta	cctgggtggc	tcgatctctc	agctgggtga	2520
ttgggaaacc	agcgatggca	ttgcgctgag	cgcggataaa	tacacctcca	gcgatccgct	2580
gtggtatgtg	accgtgaccc	tgccggcggg	tgaatcgttt	gaatacaaat	ttatccgcac	2640
tgaaagcgat	gattccgtgg	aatgggaaag	cgatccgaac	cgcgaataca	ccgtgccgca	2700
ggcgtgcggc	acctcgacgc	cgaccgtgac	cgatacctgg	cgcggatccg	agctcaagct	2760
ttctagactc	gaggaattcg	gtaccgtcga	catcgatgat	aagctgtcaa	acatgagcag	2820
atctgagccc	gcctaattgag	cgggcttttt	tttcagatct	gcttgaagac	gaaagggcct	2880
cgtgatacgc	ctatttttat	aggttaatgt	catgataata	atggtttctt	agacgatgcg	2940
tcaaagcaac	catagtagcg	gccctgtagc	ggcgcattaa	gcgcggcggg	tgtgggtggt	3000
acgcgcagcg	tgaccgctac	acttgccagc	gccctagcgc	ccgctccttt	cgtctttctc	3060
ccttcctttc	tcgccacggt	cgccggcttt	cccgcgtaag	ctctaaatcg	ggggctccct	3120
ttagggttcc	gatttagagc	tttacggcac	ctcgacccca	aaaaacttga	tttgggtgat	3180
ggttcacgta	tggggccatc	gccctgatag	acgggttttc	gccctttgac	gttgaggatcc	3240
acgttcttta	atagtggaac	ctgtttccaa	actggaacaa	cactcaaccc	tatctcgggc	3300
tattcttttg	atttataagg	gattttgcgc	atttcggcct	attgggtaaa	aatgagctg	3360
atttaacaaa	aatttaacgc	gaattttaac	aaaatattaa	cgtttacaat	ttcatcgcca	3420

ggtggcactt	ttcggggaaa	tgtgcgcgga	acccctat	gtttat	ctaaatacat	3480
tcaaatatgt	atccgtcat	gagacaataa	ccctgataa	tgcttcaata	atattgaaaa	3540
aggaagagta	gtactttaga	aaaactcatc	gagcatcaaa	tgaaaactgca	attttatcat	3600
atcaggatta	tcaataccat	atTTTTgaaa	aagccgtt	tgtaaatgaag	gagaaaactc	3660
accgaggcag	ttccatagga	tggaagatc	ctggtatcgg	tctgcgattc	cgactcgtcc	3720
aacatcaata	caacctatta	atttccctc	gtcaaaaata	agggtatcaa	gtgagaaatc	3780
accatgagtg	acgactgaat	ccggtgagaa	tggaatagc	ttatgcattt	ctttccagac	3840
ttgttcaaca	ggccagccat	tacgctcgtc	atcaaaatca	ctcgcatcaa	ccaaacggtt	3900
attcattcgt	gattgcgcct	gagcgagacg	aaatacgcga	tgcgtgttaa	aaggacaatt	3960
acaaacagga	atcgaatgca	accggcgcag	gaacactgcc	agcgcatcaa	caatatTTtc	4020
acctgaatca	ggatattctt	ctaatacctg	gaatgctgtt	ttcccgggga	tgcagtggtt	4080
gagtaaccat	gcatacatcg	gagtacggat	aaaatgcttg	atggtcggaa	gaggcataaa	4140
ttccgtcagc	cagtttagtc	tgacctctc	atctgtaaca	tcattggcaa	cgctaccttt	4200
gccaatgttc	agaaacaact	ctggcgcatc	gggcttccca	tacagtcgat	agattgtcgc	4260
acctgattgc	ccgacattat	cgcgagccca	tttataccca	tataaatcag	catccatgtt	4320
ggaatttaat	cgcggcctgg	agcaagacgt	ttcccgttga	atatggctca	taacacccct	4380
tgtattactg	tttatgtaag	cagacagttt	tattgttcat	gatgatatat	ttttatcttg	4440
tgcaatgtaa	catcagagat	tttgagacac	aacgtggctt	ttcccccccc	cccgatcgga	4500
ggaccgaagg	agctaaccgc	ttttttgcac	aacatggggg	atcatgtaac	tgccttgat	4560
cgttgggaac	cggagctgaa	tgaagccata	ccaaacgacg	agcgtgacac	cacgatgcct	4620
gcagcaatgg	caacaacggt	gcgcaaaacta	ttactggcg	aactacttac	tctagcttcc	4680
cggcaacaat	taatagactg	gatggaggcg	gataaagttg	caggaccact	tctgcgctcg	4740
gcccttccgg	ctggctggtt	tattgctgat	aaatctggag	ccggtgagcg	tgggtctcgc	4800
ggtatcattg	cagcactggg	gccagatggt	aagccctccc	gtatcgtagt	tatctacacg	4860
acggggagtc	aggcaactat	ggatgaacga	aatagacaga	tgcgtgagat	agggtcctca	4920
ctgattaagc	attggttaact	gtcagaccaa	gtttactcat	atatacttta	gattgattta	4980
aaacttcatt	tttaatttaa	aaggatctag	gtgaagatcc	tttttgataa	tctcatgacc	5040
aaaatccctt	aacgtgagtt	ttcgttccac	tgagcgtcag	accccgtaga	aaagatcaaa	5100
ggatcttctt	gagatccctt	ttttctgcgc	gtaatctgct	gcttgcaaac	aaaaaaacca	5160
ccgctaccag	ccggtggttg	tttgccggat	caagagctac	caactctttt	tccgaaggta	5220
actggcttca	gcagagcgca	gataccaaat	actgtccttc	tagtgtagcc	gtagttaggc	5280

caccacttca agaactctgt agcaccgcct acatacctcg ctctgctaata cctgttacc 5340
 gtggctgctg ccagtgccga taagtctgtg cttaccgggt tggactcaag acgatagtta 5400
 ccggataagg cgcagcggtc gggctgaacg ggggggttcgt gcacacagcc cagcttgagg 5460
 cgaacgacct acaccgaact gagataccta cagcgtgagc tatgagaaag cgccacgctt 5520
 ccgaaggga 'gaaaggcgga cagggtatccg gtaagcggca gggctcggaac aggagagcgc 5580
 acgagggagc ttccaggggg aaacgcctcg tatctttata gtcctgtcgg gtttcgccac 5640
 ctctgacttg agcgtcgatt tttgtgatgc tcgtcagggg ggcggagcct atggaaaaac 5700
 gccagcaacg cggccttttt acgggttcctg gccttttgct ggccttttgc tcacatgttc 5760
 ttctctcgt tatccctga ttctgtggat aaccgtatta ccgcctttga gtgagctgat 5820
 accgctcgcc gcagccgaac gaccgagcgc agcgagtcag tgagcgagga agcgggaagag 5880
 cgctctgatgc ggtattttct ccttacgcat ctgtgcggta tttcacaccg cataatggtg 5940
 cactctcagt acaatctgct ctgatgccgc atagttaagc cagtatacac tccgctatcg 6000
 ctacgtgact gggctcatggc tgcgccccga cccccccaa caccgctga cgcgccctga 6060
 cgggcttctc tgctcccgcc atccgcttac agacaagctg tgaccgtctc cgggagctgc 6120
 atgtgtcaga ggttttcacc gtcacaccg aaacgcata cgagccggaa gcaataaagt 6180
 taaagcctgg ggtgcctaata gagttagcta acttacatta attgctgtgc gctcactgcc 6240
 cgctttccag tcgggaaacc tgctgtgcca gctgcattaa tgaatcgcc aacgcgcggg 6300
 gagagggcgt ttgcgtattg ggcgccaggg tggtttttct ttccaccagt gagacgggca 6360
 acagctgatt gcccttcacc gcctggccct gagagagttg cagcaagcgg tccacgctgg 6420
 ttgccccag caggcgaaaa tcctgtttga tgggtggtta cggcgggata taacatgagc 6480
 tgtcttcggt atcgtcgtat cccactaccg agatatccgc accaaccgcg agcccgagct 6540
 cggtaatggc gcgcattcgc ccagcgcca tctgatcggt ggcaaccagc atcgcagtg 6600
 gaacgatgcc ctcatctcagc atttgcattg tttgttgaaa accggacatg gcactccag 6659

<210> 11

<211> 6647

<212> DNA

<213> Artificial

<220>

<223> Custom DNA vector pMXS39

<400> 11

tcttttcacc agtgagacgg gcaacagctg attgcccttc accgcctggc cctgagagag 60

ttgcagcaag cgggtccacg tgggttggcc cagcaggcga aaatcctgtt tgatggtggt 120

taaaggcggg atataacatg agctgtcttc ggtatcgtcg tatccacta ccgagatata 180

cgcaccaacg cgcagcccg	actcggtaat ggcgcgcatt	gcgccacgc ccactctgac	240
gttggcaacc agcatcgacg	tgggaacgat gccctcatte	agcatttgca tgggttgttg	300
aaaaccggac atggcactcc	agtcgccttc ccgttcgcct	atcggctgaa tttgatlgcg	360
agtgcagatc ttatgccagc	cagccagacg cagacgcgcc	gagacagaac ttaattgggcc	420
cgctaacagc gcgatttgct	ggtgacccaa tgcgaccaga	tgctccacgc ccagtcgcgt	480
accgtcttca tgggagaaaa	taatactggt gatgggtgtc	tggtcagaga catcaagaaa	540
taacgccgga acattagatg	aggcagcttc cacagcaatg	gcacccctgt catccagcgg	600
atagttaatg atcagccacc	tgacgcgttg cgcgagaaga	ttgtgcaccg ccgctttaca	660
ggcttcgacg ccgcttcggt	ctaccatcga caccaccacg	ctggcaccga gttgatcggc	720
gcgagattta atcgccgcga	caatttgcca cggcgcgtgc	agggccagac tggaggtggc	780
aacgccaatc agcaacgact	gtttgccgcg cagttgttgt	gccacgcggt tgggaatgta	840
atlcagctcc gccatcgccg	cttcacattt tccccgcgtt	ttcgcagaaa cgtggctggc	900
ctggttcacc acgcgggaaa	cggctcgata agagacacgc	gcatactctg cgacatcgta	960
taacgttact ggtttcacat	tcaccaccct gaattgactc	tctccgggga gctatcatgc	1020
cataccgcga aagggttttc	accattcgat ggtgtccttg	cacgacaggt tccccgactg	1080
gaaagcgggc agtgagcgca	acgcaattaa tgtaagttag	ctcactcatt aggcacccca	1140
ggctttacac tttatgcttc	cggctcgatc aatgtgtgga	attgtgagcg gataacaatt	1200
tcacacagga aacaggatcg	atccatcgat gagcttactc	cccatcccc tggtgacaat	1260
taatcatcgg ctcgataaat	gtgtggaatt gtgagcggat	aacaatttca cacaggaac	1320
aggatcagct tactcccat	ccccctgttg acaattaatc	atcggctcgt ataattgttg	1380
gaattgtgag cggataacaa	tttcacacag gaaacaggat	ctatcgatgc ttaggaggtc	1440
atatgaaat cgaagaaggt	aaactggtaa tctggattaa	cggcgataaa ggctataacg	1500
gtctcgtcga agtcggtaag	aaattcgaga aagataccgg	aattaaagtc accgttgagc	1560
atccggataa actggaagag	aaattccac aggttgcggc	aactggcgat gcccttgaca	1620
ttatcttctg ggcacacgac	cgctttggtg gctacgctca	atctggcctg ttggctgaaa	1680
tcaccccgga caaagcgctc	caggacaagc tgtatccgtt	tacotgggat gccgtacgtt	1740
acaacggcaa gctgattgct	taccgcgacg ctgttggaagc	gttatcgctg atttataaca	1800
aagatctgct gccgaacccg	ccaaaaacct ggggaagagat	ccggcgctg gataaagaac	1860
tgaagcgaa aggttaagagc	gcgctgatgt tcaacctgca	agaacgttac ttcacctggc	1920
cgctgattgc tgctgacggg	ggttatcgct tcaagtatga	aaacggcaag tacgacatta	1980

aagacgtggg	cgtggataac	gctggcgcca	aagcgggtct	gaccttcctg	gttgacctga	2040
ttaaaaaaa	acacatgaat	gcagacaccg	attactccat	cgcagaagct	gcctttaata	2100
aaggcgaaac	agcgatgacc	atcaacggcc	cgtgggcatg	gtccaacatc	gacaccagca	2160
aagtgaatta	tgtgttaacg	gtactgccga	ccttcaagg	tcaaccatcc	aaaccgttcg	2220
ttggcgtgct	gagcgcaggt	attaacggcg	ccagtccgaa	caaagagctg	gcaaaagagt	2280
tcctcgaaaa	ctatctgctg	actgatgaag	gtctggaagc	ggtaataaaa	gacaaaccgc	2340
tgggtgccgt	agcgctgaag	tcttacgagg	aagagttggc	gaaagatcca	cgtattgccg	2400
ccactatgga	aaacgcccag	aaagtgaaa	tcatgccgaa	catcccgcat	atgtccgctt	2460
tctggtatgc	cgtgcgtact	gcggtgatca	acgccgccag	cggtcgctag	actgtcgatg	2520
aagccctgaa	agacgcgcag	actaattcga	gctcgaacaa	caacaacaat	aacaataaca	2580
acaacctcgg	gatcgaggga	aggatttcag	aattcggatc	cgagctcaag	ctttctagac	2640
tcgagattgt	ggcgaccggc	ggcaccacca	ccaccgcgac	cccgaccggc	tccggcagcg	2700
tgacctcgac	cagcaaaaac	accgcgaccg	cgagcaaaac	cagcaccagc	acctcatcaa	2760
cctctgttac	cacccccgac	gcggtggcgg	tgaccttcga	tctgaccgcg	accaccacct	2820
acggcgaaaa	catctacctg	gtgggctcga	tctctcagct	gggtgatttg	gaaaccagcg	2880
atggcattgc	gctgagcgcg	gataaataca	cctccagcga	tccgctgtgg	tatgtgaccg	2940
tgacctgcc	ggcgggtgaa	tcgtttgat	acaatttat	ccgattgaa	agcgatgatt	3000
ccgtggaatg	ggaaagcgat	ccgaaccgag	aatacacctg	gccgcaggcg	tgccgcacct	3060
cgaccgcgac	cgtgaccgat	acctggcgct	aatgagtcga	catcgatgat	aagctgtcaa	3120
acatgagcag	atctgagccc	gcctaagtga	cgggcttttt	tttcagatct	gcttgaagac	3180
gaaagggcct	cgtgatacgc	ctatttttat	aggtaaatgt	catgataata	atggtttctt	3240
agacgatcgc	tcaaagcaac	catagtacgc	gcctgtagc	ggcgatttaa	gcgcggcggg	3300
tgtggtggtt	acgcgcagcg	tgaccgttac	acttgccagc	gcctagcgc	ccgtcccttt	3360
cgtttttctt	ccttcctttt	tcgccacgtt	cgcggccttt	cccgctcaag	ctctaaatcg	3420
ggggctccct	ttaggggttc	gatttagagc	tttacggcac	ctcgacccca	aaaaacttga	3480
tttgggtgat	ggttcacgta	gtgggccatc	gccctgatag	acggtttttc	gccctttgac	3540
gttgagatcc	acgttcttta	atagtggact	cttgttccaa	actggaacaa	caactcaacc	3600
tatctcgggc	tattcttttg	atttataagg	gattttgcgc	atttcggcct	attggttaaa	3660
aatgagctg	atttaacaaa	aatttaacgc	gaattttaac	aaaatattaa	cgtttacaat	3720
ttcatcgta	ggtggcactt	ttcggggaaa	tgtgcgcgga	acccctattt	gtttattttt	3780
ctaaatacat	tcaaatatgt	atccgctcat	gagacaataa	ccctgataaa	tgcttcaata	3840

atattgaaaa aggaagagta gtactttaga aaaactcadc gagcatcaaa tgaaactgca	3900
atattatcat atcaggatta tcaataccat atttttgaaa aagccgtttc tgtaatgaag	3960
gagaaaaac accgaggcag ttccatagga tggcaagatc ctggtatcgg tctgcgattc	4020
cgactcgtec aacatcaata caacctatta atttccctc gtcaaaaata aggttatcaa	4080
gtgagaaac accatgagtg acgactgaat ccggtgagaa tggcaatagc ttatgcattt	4140
ctttccagac ttgttcaaca ggccagccat tacgctcgtc atcaaaatca ctgcgcatca	4200
ccaaaccgtt attcattcgt gattgcgcct gagcgagacg aaatacgcga tcgctgttaa	4260
aaggacaatt acaaacagga atcgaatgca accggcgagc gaacactgcc agcgcatcaa	4320
caatatcttc acctgaatca ggatattctt ctaatacctg gaatgctgtt ttccggggga	4380
tcgcagtggt gagtaacct gcacatcagc gactacggat aaaatgcttg atggtcggaa	4440
gaggcataaa ttccgtcagc cagttagtc tgaccatctc atctgtaaca tcattggcaa	4500
cgctaccttt gccatgtttc agaacaacct ctggcgcatc gggcttccca tacagtcgat	4560
agattgtcgc acctgattgc ccgacattat cgcgagccca ttatatacca tataaatcag	4620
catccatggt ggaatttaac cgcggcctgg agcaagacgt ttcccggtga atatggctca	4680
taacaccctt tgtattactg ttatgtgaag cagacagttt tattgttcat gatgatatat	4740
ttttatcttg tgcaatgtaa catcagagat ttgagacac aacgtggctt tcccccccc	4800
cccgatcgga ggaccgaagg agctaaccgc ttttttgcac aacatggggg atcatgtaac	4860
tcgccttgat cgttgggaac cggagctgaa tgaagccata ccaaacgagc agcgtgacac	4920
cacgatgcct gcagcaatgg caacaacgtt gcgcaaaata ttaactggcg aactacttac	4980
tctagcttcc cggcaacaat taatagactg gatggaggcg gataaagttg caggaccact	5040
tctgcgctcg gcccttcocg ctggctgggt tattgctgat aaatctggag ccggtgagcg	5100
tgggtctcgc ggtatcattg cagcactggg gccagatggt aagccctccc gtatcgtagt	5160
tatctacacg acgggggagtc aggcactat ggatgaacga aatagacaga tcgctgagat	5220
aggtgcctca ctgattaagc attggtaact gtcagaccaa gtttactcat atatacttta	5280
gattgattta aaacttcatt tttaatttaa aaggatctag gtgaagatcc tttttgataa	5340
tctcatgacc aaaatccctt aacgtgagtt ttogttccac tgagcgtcag accccgtaga	5400
aaagatcaaa ggatcttctt gagatccttt tttctgcgc gtaatctgct gcttgcaaac	5460
aaaaaaacca ccgctaccag cgggtggttg tttccggat caagagctac caactctttt	5520
tcogaaggtg actggttcca gcagagcgca gataccaaat actgtccttc tagttagacc	5580
gtagtttagc caccacttca agaactctgt agcacgcct acatacctcg ctctgctaat	5640

```

cctgttacca gtggctgctg ccagtgccga taagtcgtgt cttaccgggt tggactcaag 5700
acgatagtta cgggataagg cgcagcggtc gggctgaacg gggggttcgt gcacacagcc 5760
cagcttgagg cgaacgacct acaccgaact gagataccta cagcgtgagc tatgagaaag 5820
cgccacgctt cccgaaggga gaaaggcgga caggtatccg gtaagcggca gggtcggaac 5880
aggagagcgc acgagggagc ttccaggggg aaacgcctgg tatctttata gtcctgtcgg 5940
gtttcgccac ctctgacttg agcgtcgatt tttgtgatgc tcgtcagggg ggcggagcct 6000
atggaaaaac gccagcaacg cggccttttt acggttcctg gcctttttgt ggccttttgc 6060
tcacatgttc tttcctgcgt tatccctga tctgtggat aacggtatta ccgcctttga 6120
gtgagctgat accgctgcc gcagccgaac gaccgagcgc agcgagtcag tgagcgagga 6180
agcgggaagag cgctgatgc ggtattttct ccttacgcat ctgtgcgga tttcacaccg 6240
cataatgggt cactctcagt acaatctgct ctgatgccgc atagttaagc cagtatacac 6300
tccgctatcg ctacgtgact gggcatggc tgcgccccga caccgcgcaa caccgcgtga 6360
cgcgccctga cgggcttgtc tgctccggc atccgcttac agacaagctg tgaccgtctc 6420
cgggagctgc atgtgtcaga ggttttcacc gtcacacccg aaacgccata cgagccggaa 6480
gcataaagtg taaagcctgg ggtgcctaata ggtgagcta acttacatta attgcgttgc 6540
gtcactgcc cgctttccag tcgggaaacc tgtcgtgccg gctgcattaa tgaatcggcc 6600
aacgcgcggg gagaggcggg ttgcgtattg ggcgccaggg tggtttt 6647

```

```

<210> 12
<211> 35
<212> DNA
<213> Artificial

```

```

<220>
<223> 5' PCR primer for pCWin2-MBP-MCS-SBD (pMXS39) expression vector

```

```

<400> 12
tgtatcctcg agattgtggc gaccgcggcg accac 35

```

```

<210> 13
<211> 38
<212> DNA
<213> Artificial

```

```

<220>
<223> 3' PCR primer for pCWin2-MBP-MCS-SBD (pMXS39) expression vector

```

```

<400> 13
aagcttgtcg actcattagc gccaggtatc ggtcacgg 38

```